

Table 4A
2006 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 1
Run Date: 5 / 15 / 07

Comparison of Racial and Ethnic Distribution by Clinic								
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	Multiple Races %	All Other / Unknown %
0101	210	98.6	0.0	0.0	0.5	0.0	1.0	0.0
0102	600	93.7	1.7	1.7	0.5	0.5	2.0	0.0
0103	652	97.9	0.6	0.5	0.0	0.0	1.1	0.0
0104	199	97.0	1.5	1.0	0.0	0.0	0.5	0.0
0250	276	90.6	0.7	0.0	1.8	0.0	6.9	0.0
0251	213	92.5	0.5	1.4	0.5	0.5	4.7	0.0
0252	216	62.0	0.9	1.9	7.4	0.5	27.3	0.0
0253	287	83.3	0.0	1.0	11.8	0.7	3.1	0.0
0401	730	97.0	0.7	0.3	0.0	0.1	1.9	0.0
0402	690	85.2	0.7	1.3	0.1	0.0	12.6	0.0
0403	262	98.5	0.0	0.0	0.0	0.0	1.5	0.0
0404	279	92.5	0.0	2.9	0.0	0.4	4.3	0.0
0501	733	95.9	0.3	0.7	0.1	0.0	3.0	0.0
0502	8	*	*	*	*	*	*	*
0503	250	93.2	0.0	3.6	0.0	0.0	3.2	0.0
0504	1,024	87.0	0.6	1.5	3.5	0.7	6.7	0.0
0505	650	89.5	0.9	3.2	0.5	0.0	5.8	0.0
0506	343	91.3	0.0	4.4	0.3	0.0	4.1	0.0
0650	1,446	83.1	2.4	5.3	0.5	1.2	7.5	0.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 4A
2006 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 2
Run Date: 5 / 15 / 07

Comparison of Racial and Ethnic Distribution by Clinic								
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	Multiple Races %	All Other / Unknown %
0651	957	95.6	0.3	1.0	0.1	0.1	2.8	0.0
0652	604	91.6	0.0	3.5	0.2	0.0	4.8	0.0
0653	732	97.5	0.0	0.3	0.1	0.3	1.8	0.0
0654	513	96.1	0.8	1.6	0.0	0.0	1.6	0.0
0655	642	96.0	0.3	0.5	0.2	0.3	2.8	0.0
0657	191	91.6	1.0	1.0	0.0	0.0	6.3	0.0
0701	636	90.7	1.3	0.5	0.5	5.2	1.9	0.0
0702	165	89.1	0.0	0.0	0.0	0.0	10.9	0.0
0703	355	88.2	2.8	1.1	0.3	0.6	7.0	0.0
0704	132	97.0	0.0	1.5	0.0	0.0	1.5	0.0
0705	328	98.2	0.3	0.0	0.3	0.0	1.2	0.0
0850	1,487	86.1	3.5	6.4	0.0	0.2	3.8	0.0
0851	1,233	91.2	0.7	4.9	0.4	0.2	2.6	0.0
0901	2,657	74.3	2.4	13.6	0.1	0.2	9.4	0.0
0902	126	95.2	0.0	3.2	0.0	0.0	1.6	0.0
1051	2,081	89.7	0.8	5.7	0.6	0.4	2.8	0.0
1101	2,414	29.7	60.2	4.6	0.2	0.7	4.6	0.0
1102	1,224	71.3	12.9	5.9	0.2	0.2	9.4	0.0
1106	174	89.1	1.7	3.4	0.0	0.0	5.7	0.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 4A
2006 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 3
Run Date: 5 / 15 / 07

Comparison of Racial and Ethnic Distribution by Clinic								
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	Multiple Races %	All Other / Unknown %
1250	1,583	84.7	1.1	12.6	0.1	0.2	1.3	0.0
1251	1,075	94.7	0.5	2.4	0.0	0.1	2.3	0.0
1252	1,263	56.6	0.7	41.3	0.1	0.1	1.3	0.0
1253	1,146	81.8	9.3	5.3	0.1	0.2	3.3	0.0
1350	3,567	56.9	23.2	11.6	0.1	0.8	7.3	0.0
1351	313	92.3	0.6	3.8	0.0	0.3	2.9	0.0
1352	730	58.5	30.8	4.8	0.0	0.1	5.8	0.0
1401	420	88.3	0.2	3.6	0.0	0.0	7.9	0.0
1402	283	47.0	0.4	28.6	12.7	0.0	11.3	0.0
1403	67	*	*	*	*	*	*	*
1404	38	*	*	*	*	*	*	*
1501	1,683	45.0	8.1	41.2	0.1	0.1	5.4	0.0
1502	186	89.8	1.1	3.8	0.5	0.5	4.3	0.0
1503	1,736	44.5	5.9	38.8	0.5	0.5	9.7	0.0
1504	3,562	35.2	3.4	53.2	0.1	3.5	4.6	0.0
1505	1,124	80.1	1.6	11.9	0.0	0.7	5.7	0.0
1506	866	74.9	1.5	18.7	0.0	0.1	4.7	0.0
1507	945	89.6	0.7	5.7	0.0	0.4	3.5	0.0
1508	402	0.5	0.0	99.5	0.0	0.0	0.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 4A
2006 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 4
Run Date: 5 / 15 / 07

Comparison of Racial and Ethnic Distribution by Clinic								
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	Multiple Races %	All Other / Unknown %
1509	1,019	49.8	2.4	46.1	0.0	0.0	1.8	0.0
1510	341	85.6	3.8	9.1	0.0	0.0	1.5	0.0
1511	765	89.3	1.7	4.8	0.0	0.0	4.2	0.0
1512	457	88.6	1.1	6.6	0.2	0.2	3.3	0.0
1701	939	51.1	0.3	1.9	19.3	0.9	26.4	0.1
1712	217	53.5	0.0	0.5	30.9	0.0	15.2	0.0
1950	1,039	94.4	0.5	4.1	0.4	0.1	0.5	0.0
1952	264	97.7	0.8	0.4	0.0	0.0	1.1	0.0
1953	515	93.2	0.0	4.9	0.0	0.0	1.9	0.0
1954	599	83.5	0.0	13.7	0.0	0.0	2.8	0.0
1955	334	82.0	0.0	13.8	0.0	0.3	3.9	0.0
1957	603	86.4	1.3	8.6	0.0	0.0	3.6	0.0
1959	314	81.2	5.1	8.9	0.0	0.0	4.8	0.0
1963	291	86.9	0.0	11.3	0.0	0.0	1.7	0.0
2150	1,103	88.6	0.1	1.2	0.6	0.1	9.4	0.0
2151	459	93.5	0.0	3.5	0.0	0.0	3.1	0.0
2251	790	95.7	0.6	0.9	0.0	0.5	2.3	0.0
2252	290	88.6	0.3	3.8	0.0	0.0	7.2	0.0
2551	4,913	28.9	65.8	2.2	0.1	0.2	2.8	0.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 4A
2006 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 5
Run Date: 5 / 15 / 07

Comparison of Racial and Ethnic Distribution by Clinic								
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	Multiple Races %	All Other / Unknown %
2552	1,236	28.1	61.9	4.3	0.0	0.1	5.7	0.0
2554	210	89.5	4.3	4.3	0.0	0.0	1.9	0.0
2555	5,557	77.5	16.6	2.7	0.1	0.5	2.5	0.0
3001	1,712	8.4	69.4	18.3	0.1	0.1	3.7	0.1
3002	409	15.2	57.0	25.4	0.0	0.0	2.4	0.0
3003	25	*	*	*	*	*	*	*
3004	450	0.7	0.0	99.3	0.0	0.0	0.0	0.0
3007	61	*	*	*	*	*	*	*
3009	21	*	*	*	*	*	*	*
3201	1,004	92.8	0.1	4.8	0.3	0.1	1.9	0.0
3301	4,790	49.7	25.1	16.1	0.1	4.0	4.9	0.0
3302	659	26.1	38.8	23.1	0.2	6.4	5.5	0.0
3303	105	95.2	1.0	2.9	0.0	0.0	1.0	0.0
3304	213	22.5	16.0	9.4	0.0	50.2	1.9	0.0
3305	170	73.5	11.2	7.1	0.0	4.1	4.1	0.0
3306	156	91.7	3.8	2.6	0.0	0.0	1.9	0.0
3308	82	*	*	*	*	*	*	*
3309	56	*	*	*	*	*	*	*
3311	10	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 4A
2006 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 6
Run Date: 5 / 15 / 07

Comparison of Racial and Ethnic Distribution by Clinic								
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	Multiple Races %	All Other / Unknown %
3550	131	2.3	0.0	2.3	9.2	0.0	86.3	0.0
3551	44	*	*	*	*	*	*	*
3552	79	*	*	*	*	*	*	*
3650	16	*	*	*	*	*	*	*
3701	1,051	79.1	0.4	15.2	0.0	0.6	4.8	0.0
3702	302	86.8	0.3	8.3	0.0	0.0	4.6	0.0
3703	154	87.7	0.0	8.4	0.0	0.0	3.9	0.0
3704	152	78.3	0.7	15.8	0.0	0.0	5.3	0.0
3805	3,950	70.4	14.4	7.2	0.3	0.3	7.4	0.0
3950	2,839	56.5	20.5	9.3	0.1	1.2	12.3	0.0
3952	101	92.1	0.0	4.0	0.0	0.0	4.0	0.0
3953	134	59.7	17.9	8.2	0.7	2.2	11.2	0.0
3954	282	74.1	6.7	8.5	0.0	1.8	8.9	0.0
4001	1,507	18.0	59.1	14.3	0.1	1.4	7.2	0.0
4002	1,229	40.4	29.9	18.8	0.1	1.3	9.4	0.0
4103	102	11.8	68.6	10.8	1.0	5.9	2.0	0.0
4106	2,454	4.3	46.4	45.8	0.1	0.3	3.1	0.0
4107	4,219	46.3	27.4	16.0	0.3	1.6	8.3	0.0
4109	1,583	82.6	0.7	9.8	0.4	0.2	6.4	0.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 4A
2006 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 7
Run Date: 5 / 15 / 07

Comparison of Racial and Ethnic Distribution by Clinic								
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	Multiple Races %	All Other / Unknown %
4110	2,499	25.3	8.1	59.6	0.4	0.1	6.4	0.0
4111	56	*	*	*	*	*	*	*
4112	1,121	0.6	0.2	99.2	0.0	0.0	0.0	0.0
4113	999	3.8	12.8	81.5	0.0	0.6	1.3	0.0
4120	2,951	55.3	9.3	27.0	0.1	1.7	6.7	0.0
4160	2,675	37.2	32.1	14.5	0.1	7.8	8.3	0.0
4351	1,084	89.4	0.9	5.0	0.3	0.1	4.3	0.0
4352	328	66.8	13.4	1.8	0.3	0.3	17.4	0.0
4353	778	69.8	0.3	24.3	0.3	0.3	5.1	0.0
4354	1,173	85.8	4.0	3.7	0.0	0.2	6.3	0.0
4355	538	81.6	0.0	5.4	0.6	0.9	11.5	0.0
4356	860	77.8	0.7	13.4	0.2	0.5	7.4	0.0
4357	12	*	*	*	*	*	*	*
4358	441	2.9	0.0	96.6	0.0	0.0	0.5	0.0
4359	560	65.0	0.0	33.0	0.0	0.2	1.8	0.0
4360	262	91.2	0.0	6.1	0.0	0.0	2.7	0.0
4361	360	90.8	1.4	3.3	0.6	0.0	3.9	0.0
4362	517	93.8	0.0	2.5	0.0	0.0	3.7	0.0
4363	1,244	91.4	0.2	2.3	0.0	0.2	5.8	0.1

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 4A
2006 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 8
Run Date: 5 / 15 / 07

Comparison of Racial and Ethnic Distribution by Clinic								
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	Multiple Races %	All Other / Unknown %
4401	1,669	88.3	0.6	7.0	0.0	0.1	4.1	0.0
4501	1,622	92.5	0.2	5.1	0.1	0.1	1.9	0.0
4601	1,973	64.1	1.8	29.1	0.1	0.3	4.6	0.0
4701	1,368	91.7	1.5	4.4	0.1	1.2	1.0	0.0
5050	1,654	71.6	16.4	2.8	0.1	1.5	7.7	0.0
5051	4,476	66.4	16.2	8.1	0.1	2.3	7.0	0.0
5052	2,587	69.3	13.1	2.6	0.2	7.0	7.7	0.0
5054	330	45.2	8.2	43.3	0.0	0.9	2.4	0.0
5202	470	88.1	0.2	1.7	0.4	0.2	9.4	0.0
5203	321	83.2	0.3	2.5	1.6	0.3	12.1	0.0
5204	214	85.0	1.4	1.9	0.5	0.0	11.2	0.0
5205	357	90.2	0.0	2.0	0.8	0.0	7.0	0.0
5601	1,581	92.2	1.0	2.0	0.2	0.1	4.5	0.0
5801	2,102	80.7	4.2	8.2	0.1	0.4	6.3	0.0
5803	26	*	*	*	*	*	*	*
6103	5,977	53.2	26.0	11.7	0.6	0.6	7.9	0.0
6301	6,211	38.8	27.5	27.8	0.1	2.1	3.7	0.0
6312	3,639	43.2	48.8	1.7	0.1	2.0	4.2	0.0
6316	1,513	75.5	8.9	9.1	0.0	0.9	5.6	0.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 4A
2006 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 9
Run Date: 5 / 15 / 07

Comparison of Racial and Ethnic Distribution by Clinic								
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	Multiple Races %	All Other / Unknown %
6320	396	68.7	22.7	1.8	1.8	2.5	2.5	0.0
6325	233	87.6	4.3	3.9	0.4	1.7	2.1	0.0
6330	377	24.4	53.3	18.3	0.5	1.1	2.4	0.0
6335	860	76.6	11.3	3.5	0.2	2.6	5.7	0.1
6340	632	50.0	13.1	30.9	0.2	3.5	2.4	0.0
7304	450	88.4	1.1	8.2	0.0	0.0	2.2	0.0
7306	3,847	40.9	35.6	18.0	0.0	0.4	5.1	0.0
7308	343	85.7	1.7	8.2	0.0	0.0	4.4	0.0
7401	3,219	76.4	5.3	7.7	0.3	0.4	9.9	0.0
7402	479	89.8	0.4	4.0	0.6	0.0	5.2	0.0
7403	127	80.3	0.0	16.5	0.0	0.0	3.1	0.0
7404	186	91.9	1.6	2.7	0.5	0.5	2.7	0.0
7601	1,124	92.7	0.0	5.2	0.0	0.0	2.0	0.0
7850	1,530	91.8	0.5	3.9	0.1	0.0	3.7	0.0
7851	373	97.3	0.0	1.3	0.0	0.0	1.3	0.0
7852	192	93.2	1.0	1.0	1.6	0.0	3.1	0.0
8150	156	50.6	34.6	7.1	0.0	3.2	4.5	0.0
8151	3,333	35.7	44.9	13.3	0.2	1.6	4.3	0.0
8152	498	30.5	45.4	7.4	0.0	0.0	16.7	0.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 4A
2006 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 10
Run Date: 5 / 15 / 07

Comparison of Racial and Ethnic Distribution by Clinic								
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	Multiple Races %	All Other / Unknown %
8153	57	*	*	*	*	*	*	*
8154	542	35.2	20.8	9.8	0.4	31.2	2.6	0.0
8155	134	64.9	23.1	4.5	0.0	2.2	5.2	0.0
8156	65	*	*	*	*	*	*	*
8157	28	*	*	*	*	*	*	*
8201	3,020	55.7	29.2	5.5	0.4	1.4	7.8	0.0
8202	4,882	87.3	4.6	5.3	0.1	1.5	1.2	0.0
8255	3,963	72.4	21.2	3.1	0.1	0.9	2.3	0.0
8257	866	40.0	26.2	0.1	0.1	31.1	2.5	0.0
8258	2,305	66.2	22.4	5.1	0.1	0.9	5.2	0.0
8901	1,414	0.4	96.7	0.6	0.2	0.4	1.6	0.0
8902	745	2.0	93.7	0.8	0.0	0.3	3.2	0.0
8903	1,744	2.1	93.8	0.7	0.1	1.5	1.9	0.0
8904	1,307	7.4	64.2	1.8	0.6	23.9	2.0	0.0
8905	2,834	16.3	27.3	37.6	0.2	12.6	6.0	0.0
9004	547	0.0	98.0	0.5	0.0	0.2	1.3	0.0
9010	248	1.2	98.8	0.0	0.0	0.0	0.0	0.0
9023	46	*	*	*	*	*	*	*
9040	473	0.2	98.1	0.4	0.0	0.0	1.3	0.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 4A
2006 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 11
Run Date: 5 / 15 / 07

Comparison of Racial and Ethnic Distribution by Clinic								
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	Multiple Races %	All Other / Unknown %
9045	1,781	7.2	45.7	45.6	0.1	0.5	1.0	0.0
9050	2,822	16.7	45.5	34.5	0.1	2.8	0.4	0.0
9051	3,425	4.1	6.0	88.7	0.1	0.1	1.0	0.0
9053	3,298	0.4	97.8	0.2	0.2	0.9	0.5	0.0
9054	746	2.0	92.1	2.1	0.1	1.3	2.3	0.0
9058	912	81.8	13.7	0.1	0.1	3.9	0.3	0.0
9066	2,943	0.5	98.8	0.3	0.0	0.2	0.2	0.0
9068	2,017	3.3	95.2	0.3	0.0	0.6	0.5	0.0
9069	2,326	1.1	97.1	0.3	0.0	0.3	1.2	0.0
9070	2,548	1.2	86.8	0.1	0.2	11.0	0.6	0.0
9072	1,963	1.0	95.1	0.1	0.0	3.6	0.2	0.0
9074	1,679	0.9	91.2	1.4	0.1	0.2	6.1	0.0
9079	1,635	0.5	98.0	0.2	0.1	0.3	1.0	0.0
9080	3,385	35.1	58.6	3.6	0.1	0.1	2.5	0.0
9151	20	*	*	*	*	*	*	*
Unknown Clinics	1,299	11.4	78.9	0.7	0.2	0.3	8.5	0.1
Michigan	232,137	53.1	26.5	13.7	0.4	1.6	4.7	0.0
Nation (Prior Year)	7,117,562	32.3	18.5	37.7	1.0	2.5	1.0	7.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5A
2006 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 1
Run Date: 5 / 15 / 07

Comparison of Age Distribution by Clinic				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
0101	210	39.5	16.7	43.8
0102	600	38.0	19.5	42.5
0103	652	34.8	21.0	44.2
0104	199	35.7	22.6	41.7
0250	276	31.9	23.2	44.9
0251	213	28.6	22.5	48.8
0252	216	31.9	22.2	45.8
0253	287	26.5	23.0	50.5
0401	730	35.8	21.6	42.6
0402	690	35.5	20.3	44.2
0403	262	30.9	19.8	49.2
0404	279	27.6	19.7	52.7
0501	733	24.0	23.9	52.1
0502	8	*	*	*
0503	250	40.0	18.0	42.0
0504	1,024	34.4	18.9	46.7
0505	650	31.8	20.3	47.8
0506	343	30.0	20.4	49.6

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5A
2006 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 2
Run Date: 5 / 15 / 07

Comparison of Age Distribution by Clinic				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
0650	1,446	38.5	21.4	40.0
0651	957	35.2	21.0	43.8
0652	604	32.5	21.2	46.4
0653	732	31.7	21.4	46.9
0654	513	36.3	21.4	42.3
0655	642	32.9	20.7	46.4
0657	191	30.9	24.6	44.5
0701	636	39.3	18.2	42.5
0702	165	33.3	21.8	44.8
0703	355	37.5	19.4	43.1
0704	132	37.9	23.5	38.6
0705	328	28.4	19.5	52.1
0850	1,487	37.5	20.9	41.6
0851	1,233	35.4	20.0	44.7
0901	2,657	36.3	19.7	44.0
0902	126	33.3	17.5	49.2
1051	2,081	34.5	20.3	45.3
1101	2,414	32.3	23.9	43.9

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5A
2006 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 3
Run Date: 5 / 15 / 07

Comparison of Age Distribution by Clinic				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
1102	1,224	38.0	22.3	39.7
1106	174	39.1	21.3	39.7
1250	1,583	36.6	19.8	43.6
1251	1,075	29.5	24.2	46.3
1252	1,263	33.1	22.9	44.0
1253	1,146	38.3	19.2	42.5
1350	3,567	37.3	20.2	42.6
1351	313	30.0	24.6	45.4
1352	730	30.5	24.2	45.2
1401	420	36.7	19.3	44.0
1402	283	35.7	22.3	42.0
1403	67	*	*	*
1404	38	*	*	*
1501	1,683	32.5	21.0	46.5
1502	186	35.5	25.3	39.2
1503	1,736	32.2	20.6	47.2
1504	3,562	36.6	21.5	41.9
1505	1,124	32.7	20.3	47.1

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5A
2006 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 4
Run Date: 5 / 15 / 07

Comparison of Age Distribution by Clinic				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
1506	866	34.1	20.8	45.2
1507	945	33.7	20.6	45.7
1508	402	29.4	23.4	47.3
1509	1,019	31.5	21.2	47.3
1510	341	30.8	24.6	44.6
1511	765	37.4	21.8	40.8
1512	457	38.9	19.3	41.8
1701	939	37.8	18.1	44.1
1712	217	31.3	16.6	52.1
1950	1,039	34.6	20.8	44.6
1952	264	31.8	26.1	42.0
1953	515	38.1	21.6	40.4
1954	599	34.7	20.0	45.2
1955	334	36.8	16.2	47.0
1957	603	35.0	19.9	45.1
1959	314	34.7	22.9	42.4
1963	291	38.8	21.6	39.5
2150	1,103	31.2	21.4	47.4

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5A
2006 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 5
Run Date: 5 / 15 / 07

Comparison of Age Distribution by Clinic				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
2151	459	34.6	18.3	47.1
2251	790	35.8	18.6	45.6
2252	290	29.0	23.4	47.6
2551	4,913	38.8	21.6	39.6
2552	1,236	44.0	17.8	38.2
2554	210	32.4	13.8	53.8
2555	5,557	41.6	19.3	39.1
3001	1,712	31.4	19.6	48.9
3002	409	29.6	19.1	51.3
3003	25	*	*	*
3004	450	28.7	14.2	57.1
3007	61	*	*	*
3009	21	*	*	*
3201	1,004	36.4	18.6	45.0
3301	4,790	29.2	23.8	47.0
3302	659	26.4	21.9	51.7
3303	105	29.5	22.9	47.6
3304	213	30.5	24.9	44.6

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5A
2006 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 6
Run Date: 5 / 15 / 07

Comparison of Age Distribution by Clinic				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
3305	170	29.4	25.9	44.7
3306	156	34.6	26.9	38.5
3308	82	*	*	*
3309	56	*	*	*
3311	10	*	*	*
3550	131	31.3	24.4	44.3
3551	44	*	*	*
3552	79	*	*	*
3650	16	*	*	*
3701	1,051	37.1	21.8	41.1
3702	302	37.4	20.9	41.7
3703	154	43.5	20.8	35.7
3704	152	39.5	19.7	40.8
3805	3,950	39.1	21.1	39.8
3950	2,839	37.2	19.5	43.4
3952	101	30.7	27.7	41.6
3953	134	38.1	14.9	47.0
3954	282	36.5	20.6	42.9

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5A
2006 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 7
Run Date: 5 / 15 / 07

Comparison of Age Distribution by Clinic				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
4001	1,507	34.2	22.4	43.4
4002	1,229	36.6	20.3	43.1
4103	102	19.6	21.6	58.8
4106	2,454	31.5	21.8	46.7
4107	4,219	33.3	21.9	44.8
4109	1,583	37.2	20.5	42.3
4110	2,499	33.1	22.0	44.9
4111	56	*	*	*
4112	1,121	34.3	21.1	44.7
4113	999	27.0	22.1	50.9
4120	2,951	36.2	21.4	42.4
4160	2,675	36.8	20.7	42.5
4351	1,084	35.2	22.1	42.6
4352	328	37.5	17.1	45.4
4353	778	32.8	21.6	45.6
4354	1,173	37.9	20.7	41.3
4355	538	33.1	22.3	44.6
4356	860	35.5	22.0	42.6

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5A
2006 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 8
Run Date: 5 / 15 / 07

Comparison of Age Distribution by Clinic				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
4357	12	*	*	*
4358	441	25.6	20.6	53.7
4359	560	33.4	21.4	45.2
4360	262	34.7	20.6	44.7
4361	360	37.2	16.7	46.1
4362	517	37.7	20.5	41.8
4363	1,244	39.2	19.9	40.9
4401	1,669	34.3	20.9	44.8
4501	1,622	33.5	19.8	46.7
4601	1,973	37.6	21.1	41.3
4701	1,368	40.1	22.7	37.2
5050	1,654	48.9	23.0	28.1
5051	4,476	46.8	23.2	30.0
5052	2,587	46.3	23.9	29.8
5054	330	43.9	24.2	31.8
5202	470	38.1	18.5	43.4
5203	321	33.0	17.1	49.8
5204	214	35.0	23.4	41.6

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5A
2006 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 9
Run Date: 5 / 15 / 07

Comparison of Age Distribution by Clinic				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
5205	357	35.9	20.4	43.7
5601	1,581	35.9	19.7	44.4
5801	2,102	50.2	21.7	28.1
5803	26	*	*	*
6103	5,977	32.4	23.1	44.5
6301	6,211	40.2	19.3	40.5
6312	3,639	40.8	19.6	39.6
6316	1,513	42.8	19.2	38.1
6320	396	36.6	23.0	40.4
6325	233	34.3	19.7	45.9
6330	377	39.5	19.9	40.6
6335	860	39.2	20.7	40.1
6340	632	41.5	24.1	34.5
7304	450	28.7	24.2	47.1
7306	3,847	33.9	22.3	43.8
7308	343	27.7	27.7	44.6
7401	3,219	37.5	20.7	41.8
7402	479	39.0	19.6	41.3

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5A
2006 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 10
Run Date: 5 / 15 / 07

Comparison of Age Distribution by Clinic				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
7403	127	42.5	23.6	33.9
7404	186	43.5	20.4	36.0
7601	1,124	33.2	20.2	46.6
7850	1,530	36.9	19.4	43.7
7851	373	27.1	22.3	50.7
7852	192	30.7	20.8	48.4
8150	156	42.3	14.7	42.9
8151	3,333	41.9	19.7	38.4
8152	498	53.8	24.1	22.1
8153	57	*	*	*
8154	542	34.7	21.8	43.5
8155	134	32.8	19.4	47.8
8156	65	*	*	*
8157	28	*	*	*
8201	3,020	47.5	20.6	31.9
8202	4,882	31.0	23.7	45.3
8255	3,963	43.0	23.0	34.1
8257	866	40.5	23.6	35.9

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5A
2006 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 11
Run Date: 5 / 15 / 07

Comparison of Age Distribution by Clinic				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
8258	2,305	37.6	22.1	40.3
8901	1,414	37.7	16.5	45.8
8902	745	36.0	18.9	45.1
8903	1,744	37.4	18.2	44.4
8904	1,307	34.3	17.7	48.0
8905	2,834	39.9	19.7	40.3
9004	547	27.4	21.9	50.6
9010	248	29.0	16.9	54.0
9023	46	*	*	*
9040	473	25.2	26.8	48.0
9045	1,781	31.0	22.9	46.1
9050	2,822	36.8	21.0	42.2
9051	3,425	32.6	22.8	44.6
9053	3,298	37.5	22.6	39.9
9054	746	48.4	23.7	27.9
9058	912	28.9	19.3	51.8
9066	2,943	36.9	24.6	38.5
9068	2,017	39.9	20.3	39.8

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5A
2006 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Comparison of Age Distribution by Clinic				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
9069	2,326	45.8	21.2	33.0
9070	2,548	39.3	22.6	38.0
9072	1,963	38.5	23.6	38.0
9074	1,679	41.8	20.5	37.6
9079	1,635	43.9	19.6	36.5
9080	3,385	29.6	20.1	50.2
9151	20	*	*	*
Unknown Clinics	1,299	36.0	18.9	45.1
Michigan	232,137	36.7	21.2	42.1
Nation (Prior Year)	7,117,562	35.3	22.0	42.6

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2006 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 1
Run Date: 5 / 15 / 07

Comparison of Growth and Anemia Indicators by Clinic (3,4)													
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)			
	Low		High	Short Stature (6)		Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
0101	45	* *	* *	206	7.8(138)	8.3(176)	18.4(172)	90	* *	* *		133	11.3 (97)
0102	140	6.4 (19)	15.0 (117)	584	6.7(106)	3.6 (88)	15.9(162)	252	21.0(147)	13.9 (93)		385	12.7 (111)
0103	144	5.6 (11)	9.0 (86)	641	7.3(126)	3.4 (80)	15.1(153)	283	21.2(151)	15.2(115)		410	6.8 (50)
0104	45	* *	* *	195	5.6 (61)	4.1(110)	11.8 (85)	79	* *	* *		124	12.9 (114)
0250	52	* *	* *	269	3.3 (3)	1.9 (18)	13.0(111)	124	14.5 (37)	16.1(124)		152	18.4 (144)
0251	40	* *	* *	213	5.2 (45)	0.0 (1)	14.1(133)	102	21.6(154)	13.7 (89)		135	10.4 (89)
0252	50	* *	* *	210	4.8 (32)	1.4 (7)	16.2(165)	97	* *	* *		123	12.2 (106)
0253	47	* *	* *	281	3.9 (10)	5.7(152)	12.8(105)	140	15.7 (68)	19.3(145)		186	16.7 (131)
0401	184	13.0 (102)	13.0 (113)	712	9.8(176)	1.4 (7)	19.9(178)	299	17.7(108)	13.0 (69)		438	10.7 (91)
0402	164	7.3 (30)	8.5 (79)	685	5.1 (41)	1.9 (18)	10.7 (57)	300	17.0 (91)	12.0 (55)		440	7.0 (52)
0403	65	* *	* *	258	4.3 (16)	9.3(182)	10.9 (63)	128	14.8 (41)	11.7 (47)		156	21.2 (150)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2006 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 2
Run Date: 5 / 15 / 07

Comparison of Growth and Anemia Indicators by Clinic (3,4)													
Clinic	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)		
	Low		High	Short Stature (6)		Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
0404	51	* *	* *	274	3.6 (6)	3.6 (88)	9.9 (36)	138	15.9 (75)	8.7 (17)		188	9.0 (75)
0501	97	* *	* *	720	3.3 (3)	5.8(154)	7.1 (4)	382	9.9 (8)	5.0 (3)		554	4.0 (17)
0502	2	* *	* *	8	* *	* *	* *	5	* *	* *		4	* *
0503	60	* *	* *	247	4.0 (12)	6.1(160)	15.0(150)	104	15.4 (62)	17.3(138)		152	4.6 (26)
0504	233	7.3 (30)	9.9 (99)	1,003	5.4 (53)	2.1 (26)	12.0 (92)	476	18.7(124)	11.8 (52)		648	3.7 (14)
0505	134	7.5 (34)	9.0 (86)	629	5.7 (65)	4.6(123)	11.3 (72)	308	17.5(101)	11.7 (47)		434	4.6 (26)
0506	66	* *	* *	339	5.0 (40)	3.5 (82)	13.9(131)	170	14.1 (33)	16.5(132)		251	1.6 (1)
0650	344	7.8 (38)	7.6 (61)	1,423	7.4(129)	3.3 (76)	10.5 (51)	568	18.7(124)	11.3 (39)		870	4.4 (22)
0651	216	8.8 (55)	6.0 (29)	931	5.2 (45)	2.9 (60)	14.1(133)	406	14.8 (41)	15.3(117)		587	12.4(108)
0652	126	6.3 (17)	12.7(111)	591	5.8 (69)	3.7 (93)	14.9(148)	267	14.6 (38)	19.9(148)		390	6.7 (49)
0653	141	6.4 (19)	8.5 (79)	710	7.7(137)	1.7 (15)	13.8(127)	339	19.5(135)	16.2(127)		490	14.3(123)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2006 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 3
Run Date: 5 / 15 / 07

Comparison of Growth and Anemia Indicators by Clinic (3,4)											
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)	
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number % (Rank)
0654	110	8.2 (43)	16.4 (118)	508	5.5 (56)	3.5 (82)	13.8 (127)	212	15.6 (67)	22.2 (152)	330 9.1 (76)
0655	125	8.8 (55)	8.8 (83)	634	7.9 (144)	3.0 (63)	12.3 (98)	290	17.6 (105)	14.1 (99)	443 6.3 (43)
0657	41	* *	* *	187	7.0 (116)	5.3 (142)	13.4 (120)	82	* *	* *	122 12.3 (107)
0701	139	2.9 (3)	8.6 (82)	628	7.0 (116)	2.7 (51)	9.1 (25)	265	18.1 (116)	9.4 (20)	357 9.8 (83)
0702	39	* *	* *	159	5.7 (65)	5.0 (132)	10.7 (57)	69	* *	* *	110 10.9 (94)
0703	89	* *	* *	350	6.3 (87)	2.3 (31)	11.4 (76)	153	16.3 (80)	14.4 (101)	201 3.0 (6)
0704	31	* *	* *	130	11.5 (184)	0.8 (3)	12.3 (98)	55	* *	* *	85 * *
0705	50	* *	* *	321	4.4 (19)	3.1 (68)	14.6 (142)	173	17.9 (113)	13.9 (93)	208 5.8 (38)
0850	344	8.4 (47)	9.6 (95)	1,445	6.8 (109)	3.5 (82)	10.1 (38)	609	19.5 (135)	13.6 (83)	903 3.7 (14)
0851	264	9.1 (64)	9.1 (90)	1,184	7.3 (126)	6.0 (158)	9.2 (26)	546	14.7 (39)	11.0 (34)	747 7.2 (54)
0901	650	8.0 (41)	7.8 (65)	2,592	7.4 (129)	3.7 (93)	10.1 (38)	1,143	17.8 (112)	13.5 (79)	1,581 7.1 (53)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2006 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 4
Run Date: 5 / 15 / 07

Comparison of Growth and Anemia Indicators by Clinic (3,4)											
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)	
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number % (Rank)
0902	24	* *	* *	123	8.9(162)	0.8 (3)	8.9 (22)	61	* *	* *	88 * *
1051	489	8.2 (43)	11.5 (107)	2,024	6.5 (95)	5.5(148)	10.1 (38)	927	15.1 (54)	9.7 (21)	1,173 9.7 (80)
1101	525	9.7 (71)	7.4 (57)	2,369	6.2 (84)	3.3 (76)	10.3 (45)	1,034	17.7(108)	13.1 (72)	1,595 20.1 (147)
1102	328	5.8 (13)	8.2 (73)	1,190	6.2 (84)	2.4 (39)	13.3(117)	474	17.7(108)	11.6 (44)	731 11.8(101)
1106	54	* *	* *	166	6.6(100)	0.6 (2)	9.6 (30)	63	* *	* *	105 15.2(129)
1250	382	7.3 (30)	7.1 (53)	1,545	7.8(138)	4.9(131)	11.7 (80)	678	12.8 (16)	10.0 (24)	1,026 12.7(111)
1251	178	2.8 (2)	7.9 (68)	1,052	4.7 (30)	5.4(146)	10.4 (49)	491	12.8 (16)	13.0 (69)	748 3.1 (8)
1252	253	6.7 (23)	4.7 (18)	1,230	6.8(109)	5.0(132)	21.6(182)	552	25.2(159)	33.3(161)	820 14.5(125)
1253	259	7.7 (36)	7.7 (63)	1,126	6.7(106)	6.0(158)	13.1(113)	490	15.3 (60)	12.4 (62)	707 10.5 (90)
1350	835	7.8 (38)	9.6 (95)	3,477	8.2(149)	3.0 (63)	12.5(101)	1,508	21.0(147)	14.0 (96)	2,239 16.8(133)
1351	48	* *	* *	304	8.2(149)	1.6 (11)	11.8 (85)	139	20.1(142)	10.1 (27)	207 12.1(104)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2006 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 5
Run Date: 5 / 15 / 07

Comparison of Growth and Anemia Indicators by Clinic (3,4)											
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)	
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number % (Rank)
1352	147	6.8 (24)	4.1 (12)	714	8.4(155)	4.2(112)	14.6(142)	331	20.8(146)	13.6 (83)	479 13.4 (116)
1401	92	* *	* *	411	4.9 (38)	5.1(135)	10.2 (43)	181	13.8 (29)	13.8 (91)	252 3.6 (10)
1402	54	* *	* *	280	4.6 (24)	2.5 (45)	18.6(173)	118	23.7(158)	24.6(155)	192 7.8 (60)
1403	8	* *	* *	65	* *	* *	* *	36	* *	* *	47 * *
1404	4	* *	* *	37	* *	* *	* *	16	* *	* *	32 * *
1501	359	7.5 (34)	7.8 (65)	1,632	5.5 (56)	4.2(112)	17.2(168)	759	17.1 (94)	17.9(141)	1,011 17.1 (134)
1502	46	* *	* *	181	4.4 (19)	6.1(160)	11.0 (65)	72	* *	* *	99 * *
1503	366	9.0 (61)	7.4 (57)	1,683	6.5 (95)	4.3(118)	13.4(120)	795	17.1 (94)	14.7(107)	1,125 15.0 (127)
1504	787	12.5 (99)	7.8 (65)	3,382	9.4(166)	2.8 (54)	13.4(120)	1,436	15.1 (54)	11.8 (52)	2,098 8.0 (65)
1505	253	9.9 (79)	11.5 (107)	1,101	5.2 (45)	4.6(123)	8.8 (19)	525	13.9 (31)	8.2 (11)	720 9.7 (80)
1506	197	6.1 (16)	12.2 (110)	840	7.9(144)	4.6(123)	19.3(177)	375	17.3 (98)	15.5(120)	552 13.8 (119)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2006 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 6
Run Date: 5 / 15 / 07

Comparison of Growth and Anemia Indicators by Clinic (3,4)													
Clinic	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)		
	Low		High	Short Stature (6)		Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
1507	178	4.5 (6)	14.6 (116)	927	5.6 (61)	4.2(112)	13.2(114)	433	18.0(115)	13.4 (76)		568	12.1 (104)
1508	23	* *	* *	347	14.7(186)	6.1(160)	24.5(186)	174	19.5(135)	29.9(160)		215	33.5 (168)
1509	175	8.0 (41)	8.0 (70)	992	7.9(144)	1.6 (11)	15.0(150)	470	15.7 (68)	13.6 (83)		669	17.5 (135)
1510	66	* *	* *	329	4.6 (24)	3.3 (76)	10.3 (45)	145	21.4(152)	11.0 (34)		229	10.9 (94)
1511	205	4.4 (5)	10.2 (101)	753	6.0 (78)	4.8(129)	11.8 (85)	307	19.5(135)	15.0(110)		458	24.7 (158)
1512	133	9.0 (61)	6.0 (29)	448	10.9(182)	4.0(107)	16.3(166)	192	13.5 (27)	19.3(145)		273	21.2 (150)
1701	240	6.3 (17)	13.3 (115)	922	7.6(135)	2.1 (26)	15.0(150)	401	19.2(130)	17.0(135)		615	8.8 (72)
1712	43	* *	* *	210	4.8 (32)	1.9 (18)	14.3(137)	112	17.0 (91)	15.2(115)		143	15.4 (130)
1950	251	9.6 (69)	4.8 (20)	1,017	6.5 (95)	5.2(137)	9.7 (32)	457	7.7 (5)	5.9 (6)		596	4.5 (25)
1952	61	* *	* *	262	8.0(147)	4.6(123)	12.2 (95)	113	15.0 (52)	12.4 (62)		168	6.5 (47)
1953	124	7.3 (30)	9.7 (98)	507	5.9 (72)	6.7(165)	8.3 (12)	207	9.7 (6)	6.3 (7)		302	4.6 (26)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2006 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 7
Run Date: 5 / 15 / 07

Comparison of Growth and Anemia Indicators by Clinic (3,4)													
Clinic	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)		
	Low		High	Short Stature (6)			Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
1954	135	5.9 (14)	9.6 (95)	583	7.2(122)	2.4 (39)	14.4(138)	256	14.8 (41)	14.5(105)		372	5.1 (32)
1955	86	* *	* *	330	6.4 (90)	2.4 (39)	13.0(111)	153	13.1 (21)	11.1 (38)		202	3.0 (6)
1957	143	7.7 (36)	8.4 (76)	592	4.6 (24)	4.6(123)	9.3 (27)	263	6.5 (4)	6.8 (8)		369	4.1 (19)
1959	64	* *	* *	305	7.5(132)	3.0 (63)	11.1 (67)	127	13.4 (24)	7.9 (10)		188	4.8 (29)
1963	66	* *	* *	289	8.3(151)	1.4 (7)	14.2(135)	112	17.9(113)	13.4 (76)		176	4.0 (17)
2150	235	9.8 (75)	8.5 (79)	1,080	5.5 (56)	3.1 (68)	11.7 (80)	503	18.7(124)	16.3(130)		694	1.9 (3)
2151	110	5.5 (10)	6.4 (40)	454	5.9 (72)	3.1 (68)	7.9 (10)	209	14.8 (41)	12.4 (62)		290	2.4 (4)
2251	187	9.1 (64)	7.0 (51)	775	5.9 (72)	1.9 (18)	12.9(107)	357	19.6(139)	16.2(127)		484	5.8 (38)
2252	52	* *	* *	286	5.6 (61)	2.4 (39)	15.7(160)	137	18.2(118)	18.2(142)		184	4.9 (30)
2551	1,268	11.7 (95)	3.6 (6)	4,833	6.9(111)	4.4(121)	9.5 (29)	1,924	15.1 (54)	12.0 (55)		2,932	24.7 (158)
2552	311	8.7 (53)	6.1 (34)	1,215	4.8 (32)	4.2(112)	10.9 (63)	457	18.8(127)	15.3(117)		671	23.1 (156)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2006 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 8
Run Date: 5 / 15 / 07

Comparison of Growth and Anemia Indicators by Clinic (3,4)													
Clinic	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)		
	Low		High	Short Stature (6)		Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
2554	37	* *	* *	207	6.3 (87)	6.3(164)	10.1 (38)	109	19.3(131)	10.1 (27)		127	12.6 (109)
2555	1,404	11.5 (94)	7.4 (57)	5,470	6.4 (90)	3.6 (88)	10.6 (55)	2,129	16.2 (78)	11.6 (44)		3,202	18.0 (142)
3001	386	15.8 (114)	4.4 (13)	1,643	6.5 (95)	3.5 (82)	15.4 (157)	836	15.3 (60)	13.6 (83)		1,085	12.8 (113)
3002	88	* *	* *	397	4.5 (22)	3.8 (99)	17.4 (169)	208	12.5 (13)	16.3 (130)		272	26.8 (164)
3003	4	* *	* *	23	* *	* *	* *	13	* *	* *		17	* *
3004	89	* *	* *	434	4.8 (32)	1.6 (11)	22.4 (184)	238	15.5 (65)	25.6 (157)		294	5.1 (32)
3007	11	* *	* *	61	* *	* *	* *	27	* *	* *		47	* *
3009	2	* *	* *	20	* *	* *	* *	13	* *	* *		19	* *
3201	243	9.5 (68)	9.9 (99)	992	4.8 (32)	3.6 (88)	11.7 (80)	445	14.8 (41)	13.5 (79)		642	10.0 (85)
3301	667	9.9 (79)	6.4 (40)	4,713	5.5 (56)	2.3 (31)	13.3 (117)	2,212	16.7 (84)	13.5 (79)		3,389	11.7 (100)
3302	84	* *	* *	647	5.7 (65)	3.7 (93)	12.7 (104)	327	18.3 (119)	16.2 (127)		476	10.7 (91)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2006 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 9
Run Date: 5 / 15 / 07

Comparison of Growth and Anemia Indicators by Clinic (3,4)											
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)	
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number % (Rank)
3303	13	* *	* *	103	5.8 (69)	2.9 (60)	8.7 (17)	49	* *	* *	73 * *
3304	24	* *	* *	211	4.7 (30)	2.8 (54)	8.5 (15)	96	* *	* *	160 10.0 (85)
3305	24	* *	* *	169	8.3(151)	1.8 (17)	7.1 (4)	74	* *	* *	125 6.4 (45)
3306	27	* *	* *	153	5.2 (45)	5.2(137)	12.4(100)	58	* *	* *	107 3.7 (14)
3308	12	* *	* *	82	* *	* *	* *	33	* *	* *	58 * *
3309	9	* *	* *	55	* *	* *	* *	25	* *	* *	41 * *
3311	4	* *	* *	10	* *	* *	* *	2	* *	* *	5 * *
3550	28	* *	* *	128	6.3 (87)	3.1 (68)	18.8(174)	60	* *	* *	76 * *
3551	11	* *	* *	43	* *	* *	* *	13	* *	* *	23 * *
3552	9	* *	* *	78	* *	* *	* *	44	* *	* *	55 * *
3650	1	* *	* *	15	* *	* *	* *	8	* *	* *	14 * *

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2006 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 10
Run Date: 5 / 15 / 07

Comparison of Growth and Anemia Indicators by Clinic (3,4)												
Clinic	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)	
	Low		High	Short Stature (6)		Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
3701	280	5.7 (12)	10.7 (105)	1,027	6.6(100)	5.5(148)	13.8(127)	431	22.0(157)	19.5(147)	662	6.6 (48)
3702	81	* *	* *	295	7.8(138)	4.4(121)	15.6(158)	127	13.4 (24)	20.5(150)	187	6.4 (45)
3703	40	* *	* *	150	6.0 (78)	6.7(165)	15.3(155)	57	* *	* *	85	* *
3704	40	* *	* *	149	6.7(106)	4.0(107)	10.7 (57)	61	* *	* *	92	* *
3805	1,064	10.2 (83)	7.0 (51)	3,824	6.9(111)	4.1(110)	10.5 (51)	1,502	15.2 (58)	12.8 (68)	1,925	8.8 (72)
3950	707	10.7 (89)	10.6 (103)	2,785	6.9(111)	6.2(163)	14.4(138)	1,220	18.5(123)	16.5(132)	1,693	22.8 (154)
3952	22	* *	* *	100	7.0(116)	3.0 (63)	11.0 (65)	44	* *	* *	60	* *
3953	32	* *	* *	131	5.3 (49)	8.4(177)	8.4 (13)	60	* *	* *	83	* *
3954	67	* *	* *	281	4.6 (24)	4.3(118)	15.3(155)	121	20.7(145)	18.2(142)	176	14.8 (126)
4001	323	11.8 (96)	3.7 (9)	1,444	6.2 (84)	5.3(142)	12.5(101)	641	17.0 (91)	13.7 (89)	1,029	23.2 (157)
4002	299	9.7 (71)	8.4 (76)	1,204	8.4(155)	6.9(171)	12.8(105)	519	17.1 (94)	13.5 (79)	802	22.2 (152)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2006 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 11
Run Date: 5 / 15 / 07

Comparison of Growth and Anemia Indicators by Clinic (3,4)											
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)	
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number % (Rank)
4103	11	* *	* *	102	2.9 (2)	5.9(156)	6.9 (3)	62	* *	* *	79 * *
4106	521	10.2 (83)	6.5 (44)	2,402	5.9 (72)	2.3 (31)	12.9(107)	1,122	17.1 (94)	16.0(123)	1,657 13.8(119)
4107	870	13.1 (103)	9.0 (86)	4,095	8.6(158)	2.3 (31)	13.6(124)	1,833	15.9 (75)	11.5 (41)	2,685 8.3 (67)
4109	411	10.5 (87)	9.0 (86)	1,555	6.4 (90)	2.5 (45)	9.4 (28)	671	14.8 (41)	11.6 (44)	991 3.6 (10)
4110	567	7.2 (28)	9.3 (91)	2,432	6.0 (78)	1.7 (15)	14.9(148)	1,099	21.5(153)	17.5(139)	1,616 5.9 (40)
4111	16	* *	* *	55	* *	* *	* *	16	* *	* *	30 * *
4112	258	5.4 (8)	11.6(109)	1,093	7.2(122)	1.9 (18)	23.3(185)	485	17.5(101)	27.8(159)	783 6.9 (51)
4113	168	4.2 (4)	6.5 (44)	986	4.5 (22)	3.7 (93)	16.0(163)	518	16.8 (86)	22.6(154)	714 11.3 (97)
4120	700	7.0 (26)	9.3 (91)	2,881	6.6(100)	2.5 (45)	12.9(107)	1,227	16.8 (86)	14.4(101)	1,847 8.5 (69)
4160	649	8.2 (43)	7.7 (63)	2,568	5.7 (65)	4.2(112)	12.1 (93)	1,087	14.1 (33)	14.0 (96)	1,578 4.3 (21)
4351	256	10.5 (87)	6.3 (38)	1,057	6.9(111)	2.8 (54)	13.8(127)	461	15.8 (72)	12.4 (62)	687 7.7 (59)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2006 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 12
Run Date: 5 / 15 / 07

Comparison of Growth and Anemia Indicators by Clinic (3,4)											
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)	
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number % (Rank)
4352	79	* *	* *	322	5.9 (72)	2.5 (45)	11.5 (77)	144	18.1(116)	11.8 (52)	208 19.2 (146)
4353	160	8.8 (55)	4.4 (13)	760	8.3(151)	3.4 (80)	12.2 (95)	345	18.8(127)	13.6 (83)	486 8.6 (70)
4354	275	12.4 (98)	13.1 (114)	1,158	4.9 (38)	3.8 (99)	8.4 (13)	470	14.0 (32)	9.8 (22)	704 11.9 (102)
4355	126	7.9 (40)	7.1 (53)	525	7.0(116)	5.7(152)	9.0 (24)	241	13.3 (23)	10.0 (24)	341 4.4 (22)
4356	223	8.5 (48)	4.0 (11)	843	5.3 (49)	3.7 (93)	7.1 (4)	365	14.2 (35)	8.2 (11)	531 8.9 (74)
4357	0	* *	* *	10	* *	* *	* *	5	* *	* *	8 * *
4358	68	* *	* *	424	6.4 (90)	1.2 (5)	21.9(183)	226	26.1(161)	27.0(158)	321 8.7 (71)
4359	132	8.3 (46)	10.6 (103)	535	10.1(178)	2.4 (39)	19.1(175)	256	17.6(105)	19.9(148)	345 13.0 (115)
4360	77	* *	* *	253	3.6 (6)	2.0 (23)	10.3 (45)	117	12.8 (16)	8.5 (15)	156 16.7 (131)
4361	95	* *	* *	350	7.4(129)	2.3 (31)	10.3 (45)	158	15.8 (72)	14.6(106)	216 7.9 (63)
4362	125	2.4 (1)	6.4 (40)	511	5.5 (56)	1.6 (11)	10.2 (43)	219	12.8 (16)	11.0 (34)	311 6.1 (42)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2006 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 13
Run Date: 5 / 15 / 07

Comparison of Growth and Anemia Indicators by Clinic (3,4)											
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)	
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number % (Rank)
4363	302	6.6 (22)	12.9 (112)	1,228	4.4 (19)	2.0 (23)	9.6 (30)	504	17.5 (101)	10.3 (30)	799 10.8 (93)
4401	375	9.1 (64)	10.7 (105)	1,591	10.2 (179)	3.8 (99)	11.7 (80)	720	15.7 (68)	11.4 (40)	1,002 13.5 (117)
4501	337	9.8 (75)	9.5 (94)	1,590	6.9 (111)	2.8 (54)	11.3 (72)	744	14.8 (41)	11.7 (47)	1,062 5.7 (37)
4601	493	10.8 (91)	8.1 (72)	1,925	5.6 (61)	3.8 (99)	14.8 (145)	798	17.5 (101)	17.0 (135)	802 11.6 (99)
4701	365	6.0 (15)	8.8 (83)	1,354	4.6 (24)	2.9 (60)	9.7 (32)	515	17.3 (98)	13.6 (83)	795 3.6 (10)
5050	511	12.5 (99)	7.2 (55)	1,624	7.8 (138)	2.5 (45)	14.8 (145)	459	16.6 (82)	15.3 (117)	853 25.4 (162)
5051	1,361	10.7 (89)	6.5 (44)	4,389	7.1 (121)	2.3 (31)	15.6 (158)	1,310	13.4 (24)	12.1 (58)	2,481 18.3 (143)
5052	819	9.6 (69)	5.3 (25)	2,546	7.3 (126)	2.1 (26)	15.1 (153)	765	13.1 (21)	15.0 (110)	1,376 14.3 (123)
5054	98	* *	* *	323	10.5 (181)	1.2 (5)	20.7 (181)	101	11.9 (11)	18.8 (144)	177 11.9 (102)
5202	112	5.4 (8)	8.9 (85)	461	7.8 (138)	2.6 (50)	11.7 (80)	200	20.0 (141)	11.0 (34)	251 8.0 (65)
5203	56	* *	* *	309	10.4 (180)	2.3 (31)	12.6 (103)	155	21.9 (155)	16.8 (134)	189 4.2 (20)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2006 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 14
Run Date: 5 / 15 / 07

Comparison of Growth and Anemia Indicators by Clinic (3,4)												
Clinic	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)	
	Low		High	Short Stature (6)		Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
5204	53	* *	* *	213	7.0(116)	2.8 (54)	10.8 (61)	90	* *	* *	123	4.9 (30)
5205	83	* *	* *	345	9.3(165)	2.3 (31)	14.8(145)	155	16.8 (86)	15.5(120)	186	5.9 (40)
5601	377	9.8 (75)	8.2 (73)	1,547	9.4(166)	2.1 (26)	20.5(180)	708	19.6(139)	17.1 (137)	921	7.4 (58)
5801	717	8.6 (49)	5.6 (27)	2,068	8.4(155)	2.7 (51)	8.7 (17)	586	15.9 (75)	10.8 (33)	992	9.4 (77)
5803	4	* *	* *	26	* *	* *	* *	16	* *	* *	17	* *
6103	1,538	8.6 (49)	6.5 (44)	5,776	6.6(100)	5.5(148)	10.7 (57)	2,586	18.4(122)	13.0 (69)	3,724	13.7 (118)
6301	1,594	8.6 (49)	6.2 (37)	6,102	6.6(100)	2.7 (51)	10.5 (51)	2,466	16.2 (78)	13.3 (73)	3,830	13.8 (119)
6312	944	9.7 (71)	9.3 (91)	3,558	4.6 (24)	6.8(169)	7.8 (9)	1,422	9.7 (6)	5.5 (5)	2,267	17.8 (141)
6316	426	10.1 (82)	7.3 (56)	1,484	4.3 (16)	3.3 (76)	7.1 (4)	561	16.8 (86)	9.3 (19)	905	7.8 (60)
6320	96	* *	* *	388	5.4 (53)	9.0(180)	8.8 (19)	158	17.7(108)	12.0 (55)	229	8.3 (67)
6325	48	* *	* *	228	4.8 (32)	2.2 (30)	10.5 (51)	108	17.6(105)	10.2 (29)	146	2.7 (5)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2006 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 15
Run Date: 5 / 15 / 07

Comparison of Growth and Anemia Indicators by Clinic (3,4)													
Clinic	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)		
	Low		High	Short Stature (6)		Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
6330	95	* *	* *	361	6.1 (82)	5.8(154)	12.2 (95)	148	12.8 (16)	14.9(109)		221	9.5 (78)
6335	182	7.1 (27)	6.0 (29)	840	3.8 (9)	5.6(151)	11.5 (77)	345	14.8 (41)	15.1(114)		533	7.3 (56)
6340	149	4.7 (7)	6.0 (29)	621	5.3 (49)	3.9(105)	10.8 (61)	211	14.7 (39)	13.3 (73)		384	7.3 (56)
7304	94	* *	* *	420	6.0 (78)	2.4 (39)	15.7(160)	213	21.1(149)	15.0(110)		316	4.4 (22)
7306	926	11.4 (93)	6.3 (38)	3,535	5.1 (41)	2.0 (23)	13.3(117)	1,675	16.7 (84)	12.7 (67)		2,584	10.3 (87)
7308	65	* *	* *	312	5.1 (41)	3.2 (73)	13.5(123)	155	21.9(155)	16.1(124)		253	3.6 (10)
7401	838	8.6 (49)	6.0 (29)	3,144	5.4 (53)	8.6(179)	10.4 (49)	1,336	14.8 (41)	10.0 (24)		2,031	9.8 (83)
7402	127	8.7 (53)	10.2(101)	472	7.6(135)	3.0 (63)	11.2 (69)	201	19.4(134)	13.9 (93)		303	7.9 (63)
7403	41	* *	* *	124	6.5 (95)	7.3(173)	12.9(107)	40	* *	* *		80	* *
7404	53	* *	* *	180	7.2(122)	9.4(183)	10.6 (55)	66	* *	* *		111	6.3 (43)
7601	235	6.4 (19)	5.5 (26)	1,108	4.2 (15)	5.2(137)	7.6 (8)	520	15.4 (62)	10.6 (31)		739	5.4 (34)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2006 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 16
Run Date: 5 / 15 / 07

Comparison of Growth and Anemia Indicators by Clinic (3,4)													
Clinic	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)		
	Low		High	Short Stature (6)		Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
7850	413	9.0 (61)	5.8 (28)	1,497	5.9 (72)	7.1(172)	8.8 (19)	662	15.4 (62)	11.5 (41)		943	7.8 (60)
7851	58	* *	* *	369	5.1 (41)	4.6(123)	9.8 (35)	190	21.1(149)	12.1 (58)		262	3.1 (8)
7852	35	* *	* *	190	5.8 (69)	3.2 (73)	11.6 (79)	97	* *	* *		124	5.6 (35)
8150	26	* *	* *	152	5.3 (49)	5.3(142)	11.2 (69)	66	* *	* *		72	* *
8151	831	9.9 (79)	7.6 (61)	3,146	6.6(100)	5.3(142)	11.8 (85)	1,261	20.1(142)	14.3(100)		1,905	26.1 (163)
8152	196	9.7 (71)	2.0 (1)	484	2.7 (1)	11.0(185)	8.9 (22)	106	11.3 (10)	12.3 (61)		244	20.5 (148)
8153	10	* *	* *	54	* *	* *	* *	21	* *	* *		36	* *
8154	132	9.1 (64)	6.1 (34)	510	3.9 (10)	5.1(135)	10.0 (37)	238	15.5 (65)	8.4 (14)		311	10.3 (87)
8155	30	* *	* *	125	4.0 (12)	4.8(129)	6.4 (2)	65	* *	* *		86	* *
8156	11	* *	* *	59	* *	* *	* *	26	* *	* *		35	* *
8157	3	* *	* *	28	* *	* *	* *	20	* *	* *		18	* *

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2006 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 17
Run Date: 5 / 15 / 07

Comparison of Growth and Anemia Indicators by Clinic (3,4)													
Clinic	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)						
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
8201	941	8.8 (55)	6.4 (40)	2,942	6.1 (82)	3.8 (99)	9.7 (32)	946	15.0 (52)	10.6 (31)		1,493	9.6 (79)
8202	952	6.9 (25)	7.9 (68)	4,778	4.1 (14)	3.7 (93)	11.8 (85)	2,173	14.9 (51)	13.3 (73)		2,975	5.6 (35)
8255	1,222	8.8 (55)	8.2 (73)	3,856	8.8(161)	3.8 (99)	11.9 (91)	1,308	19.3(131)	14.4(101)		1,943	7.2 (54)
8257	235	12.3 (97)	6.8 (48)	772	10.0(177)	5.2(137)	14.5(140)	300	12.7 (14)	11.7 (47)		496	25.0 (160)
8258	664	8.9 (60)	8.0 (70)	2,253	7.5(132)	2.8 (54)	13.2(114)	923	19.3(131)	16.1(124)		1,414	17.5 (135)
8901	355	14.6 (108)	4.5 (15)	1,364	9.7(173)	7.5(174)	13.6(124)	641	15.1 (54)	11.7 (47)		74	* *
8902	164	13.4 (104)	4.9 (22)	720	9.7(173)	6.7(165)	12.1 (93)	330	12.7 (14)	12.4 (62)		184	22.3 (153)
8903	415	15.7 (112)	3.6 (6)	1,681	9.0(163)	5.2(137)	14.7(144)	767	15.8 (72)	11.5 (41)		350	12.6 (109)
8904	252	13.9 (105)	2.8 (2)	1,246	7.5(132)	8.4(177)	13.9(131)	614	13.5 (27)	12.2 (60)		715	22.8 (154)
8905	690	10.3 (85)	8.4 (76)	2,685	8.3(151)	7.7(175)	16.1(164)	1,122	15.7 (68)	17.6(140)		47	* *
9004	85	* *	* *	530	9.4(166)	4.3(118)	20.2(179)	278	25.2(159)	24.8(156)		382	30.4 (167)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2006 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 18
Run Date: 5 / 15 / 07

Comparison of Growth and Anemia Indicators by Clinic (3,4)											
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)	
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number % (Rank)
9010	35	* *	* *	237	8.0(147)	4.2(112)	17.7(170)	133	20.3(144)	15.0(110)	141 39.0(171)
9023	5	* *	* *	45	* *	* *	* *	25	* *	* *	25 * *
9040	46	* *	* *	461	3.3 (3)	3.5 (82)	11.1 (67)	218	13.8 (29)	8.7 (17)	354 34.2(169)
9045	389	10.3 (85)	6.9 (49)	1,675	9.5(172)	6.7(165)	18.3(171)	780	18.3(119)	21.2(151)	177 1.7 (2)
9050	664	9.8 (75)	5.1 (24)	2,756	7.2(122)	3.5 (82)	14.2(135)	1,161	16.8 (86)	14.0 (96)	1,575 14.2(122)
9051	773	7.2 (28)	7.4 (57)	3,334	6.4 (90)	1.4 (7)	19.2(176)	1,494	18.3(119)	22.4(153)	2,034 15.0(127)
9053	882	14.5(106)	4.9 (22)	3,199	9.4(166)	6.8(169)	11.2 (69)	1,283	10.0 (9)	8.3 (13)	2,045 17.7(138)
9054	222	14.9(110)	4.5 (15)	727	9.4(166)	5.0(132)	13.6(124)	202	17.3 (98)	14.4(101)	429 36.1(170)
9058	176	10.8 (91)	3.4 (3)	815	3.7 (8)	25.8(186)	4.8 (1)	399	1.3 (1)	1.8 (1)	619 18.6(145)
9066	771	16.0(116)	3.8 (10)	2,852	11.9(185)	5.9(156)	10.1 (38)	1,111	11.9 (11)	8.6 (16)	1,213 17.7(138)
9068	544	14.5(106)	6.1 (34)	1,948	7.8(138)	9.5(184)	8.5 (15)	778	6.4 (3)	3.6 (2)	1,000 25.0(160)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2006 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 19
Run Date: 5 / 15 / 07

Comparison of Growth and Anemia Indicators by Clinic (3,4)											
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)	
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number % (Rank)
9069	669	14.9 (110)	4.6 (17)	2,260	8.6(158)	4.0(107)	11.8 (85)	759	18.8(127)	14.8(108)	1,196 17.7 (138)
9070	643	14.6 (108)	3.4 (3)	2,473	9.0(163)	9.1(181)	11.3 (72)	936	6.0 (2)	5.2 (4)	1,414 11.0 (96)
9072	548	16.6 (118)	3.6 (6)	1,918	9.4(166)	3.9(105)	11.3 (72)	727	14.4 (36)	9.9 (23)	874 9.7 (80)
9074	473	15.9 (115)	4.7 (18)	1,631	8.6(158)	3.1 (68)	13.2(114)	636	14.8 (41)	13.4 (76)	989 17.5 (135)
9079	458	15.7 (112)	3.5 (5)	1,564	9.7(173)	3.6 (88)	14.5(140)	567	16.6 (82)	13.8 (91)	856 30.0 (166)
9080	699	12.9 (101)	6.9 (49)	3,328	4.3 (16)	3.2 (73)	8.0 (11)	1,676	15.2 (58)	7.8 (9)	2,176 20.9 (149)
9151	7	* *	* *	19	* *	* *	* *	5	* *	* *	10 * *
Unknown Clinics	230	16.5 (117)	4.8 (20)	1,174	11.0(183)	5.4(146)	17.0(167)	546	16.5 (81)	15.9(122)	674 28.9 (165)
Michigan	55,559	9.8	7.3	225,665	6.8	4.1	12.4	95,780	16.2	13.3	135,168 12.9
Nation (Prior Year)	1,641,264	9.4	6.6	6,709,486	6.4	4.8	13.9	2,869,075	16.1	14.7	4,381,087 13.6

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2006 Pediatric Nutrition Surveillance (1)
Michigan

Page: 1
Run Date: 5 / 15 / 07

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
(4,5)

	Breastfeeding (5)			Exclusive Breastfeeding (6)		≤2 Hours TV Viewing/Day (7)	Smoking in Household
Clinic	Number % Ever Breastfed (Rank)	Number % Breastfed At Least 6 Months (Rank)	Number % Breastfed At Least 12 Months (Rank)	Number % Breastfed At Least 3 Months (Rank)	Number % Breastfed At Least 6 Months (Rank)	Number % (Rank)	Number % (Rank)
0101	51 * *	42 * *	21 * *	0 * *	0 * *	0 * *	0 * *
0102	150 44.7 (98)	126 8.7 (100)	73 * *	0 * *	0 * *	0 * *	0 * *
0103	154 41.6(103)	109 7.3 (105)	88 * *	0 * *	0 * *	0 * *	0 * *
0104	54 * *	38 * *	34 * *	0 * *	0 * *	0 * *	0 * *
0250	55 * *	50 * *	60 * *	0 * *	0 * *	0 * *	0 * *
0251	41 * *	35 * *	36 * *	0 * *	0 * *	0 * *	0 * *
0252	51 * *	43 * *	39 * *	0 * *	0 * *	0 * *	0 * *
0253	48 * *	52 * *	52 * *	0 * *	0 * *	0 * *	0 * *
0401	191 50.8 (75)	138 16.7 (40)	119 10.9 (56)	0 * *	0 * *	0 * *	0 * *
0402	167 69.5 (8)	121 13.2 (68)	109 10.1 (65)	0 * *	0 * *	0 * *	0 * *
0403	66 * *	41 * *	34 * *	0 * *	0 * *	0 * *	0 * *
0404	52 * *	45 * *	48 * *	0 * *	0 * *	0 * *	0 * *
0501	100 55.0 (61)	128 23.4 (16)	135 18.5 (13)	0 * *	0 * *	0 * *	0 * *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in

(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2006 Pediatric Nutrition Surveillance (1)
Michigan

Page: 2
Run Date: 5 / 15 / 07

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
(4,5)

	Breastfeeding (5)			Exclusive Breastfeeding (6)		≤2 Hours TV Viewing/Day (7)	Smoking in Household
Clinic	Number % Ever Breastfed (Rank)	Number % Breastfed At Least 6 Months (Rank)	Number % Breastfed At Least 12 Months (Rank)	Number % Breastfed At Least 3 Months (Rank)	Number % Breastfed At Least 6 Months (Rank)	Number % (Rank)	Number % (Rank)
0502	2 * *	1 * *	1 * *	0 * *	0 * *	0 * *	0 * *
0503	69 * *	47 * *	37 * *	0 * *	0 * *	0 * *	0 * *
0504	247 72.5 (7)	193 34.7 (3)	152 20.4 (9)	0 * *	0 * *	0 * *	0 * *
0505	146 60.3 (34)	104 18.3 (30)	85 * *	0 * *	0 * *	0 * *	0 * *
0506	71 * *	54 * *	62 * *	0 * *	0 * *	0 * *	0 * *
0650	373 69.4 (9)	331 17.8 (34)	251 17.9 (17)	0 * *	0 * *	0 * *	0 * *
0651	227 46.3 (92)	179 11.7 (83)	161 12.4 (38)	0 * *	0 * *	0 * *	0 * *
0652	132 53.0 (68)	102 18.6 (27)	91 * *	0 * *	0 * *	0 * *	0 * *
0653	153 59.5 (37)	142 13.4 (65)	121 9.9 (67)	0 * *	0 * *	0 * *	0 * *
0654	120 55.8 (55)	94 * *	84 * *	0 * *	0 * *	0 * *	0 * *
0655	126 55.6 (57)	143 10.5 (94)	120 9.2 (75)	0 * *	0 * *	0 * *	0 * *
0657	42 * *	35 * *	36 * *	0 * *	0 * *	0 * *	0 * *
0701	164 73.8 (6)	105 34.3 (4)	79 * *	0 * *	0 * *	0 * *	0 * *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in

(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2006 Pediatric Nutrition Surveillance (1)
Michigan

Page: 3
Run Date: 5 / 15 / 07

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
(4,5)

	Breastfeeding (5)			Exclusive Breastfeeding (6)		≤2 Hours TV Viewing/Day (7)	Smoking in Household
Clinic	Number % Ever Breastfed (Rank)	Number % Breastfed At Least 6 Months (Rank)	Number % Breastfed At Least 12 Months (Rank)	Number % Breastfed At Least 3 Months (Rank)	Number % Breastfed At Least 6 Months (Rank)	Number % (Rank)	Number % (Rank)
0702	40 * *	30 * *	31 * *	0 * *	0 * *	0 * *	0 * *
0703	89 * *	61 * *	48 * *	0 * *	0 * *	0 * *	0 * *
0704	31 * *	27 * *	24 * *	0 * *	0 * *	0 * *	0 * *
0705	60 * *	46 * *	39 * *	0 * *	0 * *	0 * *	0 * *
0850	377 63.7 (17)	280 15.4 (48)	248 14.5 (27)	0 * *	0 * *	0 * *	0 * *
0851	279 47.7 (85)	228 17.1 (37)	186 14.5 (27)	0 * *	0 * *	0 * *	0 * *
0901	665 50.1 (79)	447 10.1 (95)	380 9.7 (69)	0 * *	0 * *	0 * *	0 * *
0902	24 * *	22 * *	19 * *	0 * *	0 * *	0 * *	0 * *
1051	505 77.6 (3)	402 32.3 (6)	351 28.5 (3)	0 * *	0 * *	0 * *	0 * *
1101	533 47.1 (88)	617 11.0 (92)	470 8.5 (81)	0 * *	0 * *	0 * *	0 * *
1102	339 50.1 (79)	285 17.9 (31)	222 17.6 (18)	0 * *	0 * *	0 * *	0 * *
1106	56 * *	45 * *	29 * *	0 * *	0 * *	0 * *	0 * *
1250	402 55.7 (56)	274 13.5 (64)	247 10.5 (61)	0 * *	0 * *	0 * *	0 * *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in

(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2006 Pediatric Nutrition Surveillance (1)
Michigan

Page: 4
Run Date: 5 / 15 / 07

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
(4,5)

	Breastfeeding (5)			Exclusive Breastfeeding (6)			≤2 Hours TV Viewing/Day (7)	Smoking in Household
Clinic	Number (Rank)	% Ever Breastfed (Rank)	Number Breastfed At Least 6 Months (Rank)	% Breastfed At Least 12 Months (Rank)	Number Breastfed At Least 3 Months (Rank)	% Breastfed At Least 6 Months (Rank)	Number (Rank)	% (Rank)
1251	184	53.3 (66)	196	12.8 (73)	174	8.6 (79)	0	* *
1252	314	56.4 (51)	195	14.9 (54)	214	11.7 (43)	0	* *
1253	283	38.5(108)	197	6.1 (113)	165	2.4 (103)	0	* *
1350	910	44.8 (97)	689	8.9 (98)	582	7.9 (84)	0	* *
1351	53	* *	67	* *	55	* *	0	* *
1352	154	36.4(111)	119	7.6 (102)	138	7.2 (91)	0	* *
1401	97	* *	76	* *	67	* *	0	* *
1402	56	* *	53	* *	47	* *	0	* *
1403	8	* *	6	* *	14	* *	0	* *
1404	8	* *	8	* *	7	* *	0	* *
1501	401	62.1 (24)	293	19.8 (24)	261	15.3 (26)	0	* *
1502	47	* *	30	* *	34	* *	0	* *
1503	385	58.7 (43)	302	21.9 (19)	290	15.5 (25)	0	* *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in

(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic^(4,5)

- (1) Reporting period is January 1 through December 31.
- (2) TV Viewing and Smoking in Household data are not reported by all contributors.
- (3) Analyses based on one record per child.
- (4) Excludes records with unknown data or errors.
- (5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
- (6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
- (7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2006 Pediatric Nutrition Surveillance (1)
Michigan

Page: 6
Run Date: 5 / 15 / 07

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
(4,5)

	Breastfeeding (5)			Exclusive Breastfeeding (6)			≤2 Hours TV Viewing/Day (7)	Smoking in Household
Clinic	Number (Rank)	% Ever Breastfed (Rank)	Number % Breastfed At Least 6 Months (Rank)	Number % Breastfed At Least 12 Months (Rank)	Number % Breastfed At Least 3 Months (Rank)	Number % Breastfed At Least 6 Months (Rank)	Number (Rank)	Number (Rank)
1953	133	59.4 (39)	105 17.1 (37)	87 * *	0 * *	0 * *	0 * *	0 * *
1954	139	66.9 (11)	106 20.8 (21)	96 * *	0 * *	0 * *	0 * *	0 * *
1955	86	* *	53 * *	42 * *	0 * *	0 * *	0 * *	0 * *
1957	143	65.7 (12)	124 16.1 (43)	91 * *	0 * *	0 * *	0 * *	0 * *
1959	67	* *	56 * *	49 * *	0 * *	0 * *	0 * *	0 * *
1963	67	* *	68 * *	56 * *	0 * *	0 * *	0 * *	0 * *
2150	240	59.6 (36)	195 13.8 (61)	187 11.8 (42)	0 * *	0 * *	0 * *	0 * *
2151	111	48.6 (82)	84 * *	76 * *	0 * *	0 * *	0 * *	0 * *
2251	190	57.9 (46)	134 17.9 (31)	126 16.7 (22)	0 * *	0 * *	0 * *	0 * *
2252	53	* *	48 * *	48 * *	0 * *	0 * *	0 * *	0 * *
2551	1,301	35.3(114)	871 6.2 (112)	752 5.1 (100)	0 * *	0 * *	0 * *	0 * *
2552	334	38.0(109)	260 8.8 (99)	167 9.6 (70)	0 * *	0 * *	0 * *	0 * *
2554	44	* *	26 * *	19 * *	0 * *	0 * *	0 * *	0 * *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in

(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2006 Pediatric Nutrition Surveillance (1)
Michigan

Page: 7
Run Date: 5 / 15 / 07

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
(4,5)

	Breastfeeding (5)			Exclusive Breastfeeding (6)			≤2 Hours TV Viewing/Day (7)	Smoking in Household
Clinic	Number (Rank)	% Ever Breastfed (Rank)	Number (Rank)	% Breastfed At Least 6 Months (Rank)	Number (Rank)	% Breastfed At Least 12 Months (Rank)	Number (Rank)	% (Rank)
2555	1,532	48.3 (83)	1,046	11.4 (88)	758	7.9 (84)	0	* *
3001	389	40.1(106)	278	6.5 (107)	247	6.1 (95)	0	* *
3002	88	* *	55	* *	56	* *	0	* *
3003	4	* *	6	* *	3	* *	0	* *
3004	89	* *	51	* *	35	* *	0	* *
3007	11	* *	9	* *	10	* *	0	* *
3009	2	* *	4	* *	2	* *	0	* *
3201	247	47.0 (89)	188	11.7 (83)	150	9.3 (74)	0	* *
3301	779	62.9 (19)	1,007	17.9 (31)	813	13.5 (32)	0	* *
3302	96	* *	129	15.5 (46)	111	19.8 (11)	0	* *
3303	15	* *	23	* *	21	* *	0	* *
3304	27	* *	55	* *	43	* *	0	* *
3305	26	* *	30	* *	33	* *	0	* *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in

(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2006 Pediatric Nutrition Surveillance (1)
Michigan

Page: 8
Run Date: 5 / 15 / 07

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
(4,5)

	Breastfeeding (5)			Exclusive Breastfeeding (6)		≤2 Hours TV Viewing/Day (7)	Smoking in Household
Clinic	Number % Ever Breastfed (Rank)	Number % Breastfed At Least 6 Months (Rank)	Number % Breastfed At Least 12 Months (Rank)	Number % Breastfed At Least 3 Months (Rank)	Number % Breastfed At Least 6 Months (Rank)	Number % (Rank)	Number % (Rank)
3306	28 * *	43 * *	37 * *	0 * *	0 * *	0 * *	0 * *
3308	13 * *	18 * *	25 * *	0 * *	0 * *	0 * *	0 * *
3309	9 * *	11 * *	10 * *	0 * *	0 * *	0 * *	0 * *
3311	5 * *	4 * *	0 * *	0 * *	0 * *	0 * *	0 * *
3550	28 * *	19 * *	24 * *	0 * *	0 * *	0 * *	0 * *
3551	11 * *	6 * *	11 * *	0 * *	0 * *	0 * *	0 * *
3552	9 * *	10 * *	11 * *	0 * *	0 * *	0 * *	0 * *
3650	1 * *	1 * *	3 * *	0 * *	0 * *	0 * *	0 * *
3701	280 50.7 (76)	182 12.6 (74)	166 11.4 (47)	0 * *	0 * *	0 * *	0 * *
3702	82 * *	58 * *	49 * *	0 * *	0 * *	0 * *	0 * *
3703	40 * *	42 * *	26 * *	0 * *	0 * *	0 * *	0 * *
3704	40 * *	30 * *	20 * *	0 * *	0 * *	0 * *	0 * *
3805	1,071 50.3 (77)	910 11.9 (79)	662 6.8 (92)	0 * *	0 * *	0 * *	0 * *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in

(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2006 Pediatric Nutrition Surveillance (1)
Michigan

Page: 9
Run Date: 5 / 15 / 07

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
(4,5)

	Breastfeeding (5)						Exclusive Breastfeeding (6)		≤2 Hours TV Viewing/Day (7)	Smoking in Household
Clinic	Number % Ever Breastfed (Rank)	Number % Breastfed At Least 6 Months (Rank)	Number % Breastfed At Least 12 Months (Rank)	Number % Breastfed At Least 3 Months (Rank)	Number % Breastfed At Least 6 Months (Rank)	Number % (Rank)	Number % (Rank)			
3950	718 59.5 (37)	526 18.4 (29)	402 14.4 (29)	0 * *	0 * *	0 * *	0 * *			
3952	22 * *	21 * *	15 * *	0 * *	0 * *	0 * *	0 * *			
3953	32 * *	22 * *	17 * *	0 * *	0 * *	0 * *	0 * *			
3954	68 * *	47 * *	39 * *	0 * *	0 * *	0 * *	0 * *			
4001	332 56.9 (50)	250 11.6 (85)	250 10.8 (59)	0 * *	0 * *	0 * *	0 * *			
4002	309 61.2 (32)	220 13.6 (63)	173 12.1 (41)	0 * *	0 * *	0 * *	0 * *			
4103	12 * *	13 * *	14 * *	0 * *	0 * *	0 * *	0 * *			
4106	529 52.0 (71)	455 11.6 (85)	426 10.3 (63)	0 * *	0 * *	0 * *	0 * *			
4107	925 54.1 (63)	823 14.5 (56)	692 12.9 (37)	0 * *	0 * *	0 * *	0 * *			
4109	414 60.9 (33)	284 16.2 (42)	276 10.9 (56)	0 * *	0 * *	0 * *	0 * *			
4110	581 57.1 (48)	481 21.0 (20)	448 13.4 (33)	0 * *	0 * *	0 * *	0 * *			
4111	16 * *	15 * *	8 * *	0 * *	0 * *	0 * *	0 * *			
4112	258 92.2 (1)	173 18.5 (28)	182 17.6 (18)	0 * *	0 * *	0 * *	0 * *			

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in

(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2006 Pediatric Nutrition Surveillance (1)
Michigan

Page: 10
Run Date: 5 / 15 / 07

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
(4,5)

	Breastfeeding (5)			Exclusive Breastfeeding (6)			≤2 Hours TV Viewing/Day (7)	Smoking in Household
Clinic	Number (Rank)	% Ever Breastfed (Rank)	Number (Rank)	% Breastfed At Least 6 Months (Rank)	Number (Rank)	% Breastfed At Least 12 Months (Rank)	Number (Rank)	% (Rank)
4113	169	74.0 (4)	185	20.0 (23)	172	17.4 (21)	0	* *
4120	705	61.7 (29)	601	15.1 (53)	504	10.9 (56)	0	* *
4160	669	55.5 (58)	544	15.4 (48)	448	11.2 (50)	0	* *
4351	262	62.2 (22)	194	13.4 (65)	179	11.2 (50)	0	* *
4352	81	* *	54	* *	44	* *	0	* *
4353	164	64.0 (14)	148	15.5 (46)	128	10.2 (64)	0	* *
4354	289	64.0 (14)	243	11.9 (79)	192	9.4 (73)	0	* *
4355	126	61.9 (27)	96	* *	94	* *	0	* *
4356	224	58.0 (45)	155	22.6 (18)	138	12.3 (40)	0	* *
4357	0	* *	2	* *	2	* *	0	* *
4358	75	* *	69	* *	68	* *	0	* *
4359	140	62.1 (24)	79	* *	89	* *	0	* *
4360	78	* *	35	* *	38	* *	0	* *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in

(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2006 Pediatric Nutrition Surveillance (1)
Michigan

Page: 11
Run Date: 5 / 15 / 07

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
(4,5)

	Breastfeeding (5)			Exclusive Breastfeeding (6)		≤2 Hours TV Viewing/Day (7)	Smoking in Household
Clinic	Number % Ever Breastfed (Rank)	Number % Breastfed At Least 6 Months (Rank)	Number % Breastfed At Least 12 Months (Rank)	Number % Breastfed At Least 3 Months (Rank)	Number % Breastfed At Least 6 Months (Rank)	Number % (Rank)	Number % (Rank)
4361	95 * *	64 * *	44 * *	0 * *	0 * *	0 * *	0 * *
4362	127 74.0 (4)	87 * *	79 * *	0 * *	0 * *	0 * *	0 * *
4363	313 62.9 (19)	250 13.2 (68)	196 6.1 (95)	0 * *	0 * *	0 * *	0 * *
4401	391 51.9 (72)	249 13.3 (67)	219 11.0 (53)	0 * *	0 * *	0 * *	0 * *
4501	344 57.6 (47)	306 11.8 (81)	234 8.5 (81)	0 * *	0 * *	0 * *	0 * *
4601	497 49.9 (81)	451 13.7 (62)	350 8.9 (78)	0 * *	0 * *	0 * *	0 * *
4701	374 65.2 (13)	259 19.7 (25)	240 10.0 (66)	0 * *	0 * *	0 * *	0 * *
5050	527 43.5 (99)	384 11.5 (87)	282 7.8 (86)	0 * *	0 * *	0 * *	0 * *
5051	1,421 48.3 (83)	1,025 11.3 (90)	780 9.5 (72)	0 * *	0 * *	0 * *	0 * *
5052	836 40.2(104)	562 9.3 (96)	443 5.6 (98)	0 * *	0 * *	0 * *	0 * *
5054	106 61.3 (30)	64 * *	53 * *	0 * *	0 * *	0 * *	0 * *
5202	122 59.0 (41)	83 * *	66 * *	0 * *	0 * *	0 * *	0 * *
5203	61 * *	53 * *	42 * *	0 * *	0 * *	0 * *	0 * *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in

(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2006 Pediatric Nutrition Surveillance (1)
Michigan

Page: 12
Run Date: 5 / 15 / 07

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
(4,5)

	Breastfeeding (5)			Exclusive Breastfeeding (6)		≤2 Hours TV Viewing/Day (7)	Smoking in Household
Clinic	Number % Ever Breastfed (Rank)	Number % Breastfed At Least 6 Months (Rank)	Number % Breastfed At Least 12 Months (Rank)	Number % Breastfed At Least 3 Months (Rank)	Number % Breastfed At Least 6 Months (Rank)	Number % (Rank)	Number % (Rank)
5204	54 * *	39 * *	35 * *	0 * *	0 * *	0 * *	0 * *
5205	85 * *	55 * *	54 * *	0 * *	0 * *	0 * *	0 * *
5601	382 61.3 (30)	305 16.1 (43)	233 12.4 (38)	0 * *	0 * *	0 * *	0 * *
5801	732 52.5 (70)	580 12.6 (74)	411 9.2 (75)	0 * *	0 * *	0 * *	0 * *
5803	4 * *	2 * *	3 * *	0 * *	0 * *	0 * *	0 * *
6103	1,562 45.3 (96)	880 12.0 (78)	1,003 8.0 (83)	0 * *	0 * *	0 * *	0 * *
6301	1,675 47.3 (86)	1,263 14.1 (58)	882 11.7 (43)	0 * *	0 * *	0 * *	0 * *
6312	983 53.3 (66)	716 17.0 (39)	497 14.1 (31)	0 * *	0 * *	0 * *	0 * *
6316	427 56.0 (54)	348 14.1 (58)	212 13.2 (34)	0 * *	0 * *	0 * *	0 * *
6320	103 54.4 (62)	77 * *	65 * *	0 * *	0 * *	0 * *	0 * *
6325	49 * *	44 * *	41 * *	0 * *	0 * *	0 * *	0 * *
6330	104 33.7(115)	71 * *	43 * *	0 * *	0 * *	0 * *	0 * *
6335	209 50.2 (78)	174 20.1 (22)	138 21.0 (8)	0 * *	0 * *	0 * *	0 * *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in

(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

(2)

(4,5)

- 

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic^(4,5)

- (1) Reporting period is January 1 through December 31.
- (2) TV Viewing and Smoking in Household data are not reported by all contributors.
- (3) Analyses based on one record per child.
- (4) Excludes records with unknown data or errors.
- (5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
- (6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
- (7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2006 Pediatric Nutrition Surveillance (1)
Michigan

Page: 15
Run Date: 5 / 15 / 07

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
(4,5)

	Breastfeeding (5)						Exclusive Breastfeeding (6)				≤2 Hours TV Viewing/Day (7)	Smoking in Household		
Clinic	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months	Number	% Breastfed At Least 3 Months	Number	% Breastfed At Least 6 Months	Number	%	Number	%
	(Rank)		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)	
8902	172	37.8(110)	130	24.6 (14)	109	23.9 (5)	0	* *	0	* *	0	* *	0	* *
8903	435	29.2(120)	302	15.2 (51)	209	10.5 (61)	0	* *	0	* *	0	* *	0	* *
8904	284	51.1 (74)	198	36.9 (2)	161	26.7 (4)	0	* *	0	* *	0	* *	0	* *
8905	743	35.7(112)	457	26.0 (11)	351	23.9 (5)	0	* *	0	* *	0	* *	0	* *
9004	88	* *	108	8.3 (101)	85	* *	0	* *	0	* *	0	* *	0	* *
9010	38	* *	47	* *	31	* *	0	* *	0	* *	0	* *	0	* *
9023	5	* *	3	* *	8	* *	0	* *	0	* *	0	* *	0	* *
9040	46	* *	102	11.8 (81)	105	11.4 (47)	0	* *	0	* *	0	* *	0	* *
9045	405	40.2(104)	253	17.8 (34)	270	15.9 (24)	0	* *	0	* *	0	* *	0	* *
9050	716	51.4 (73)	592	24.3 (15)	440	21.6 (7)	0	* *	0	* *	0	* *	0	* *
9051	906	55.5 (58)	307	23.1 (17)	522	18.2 (15)	0	* *	0	* *	0	* *	0	* *
9053	961	31.0(117)	422	14.9 (54)	520	11.7 (43)	0	* *	0	* *	0	* *	0	* *
9054	225	39.1(107)	173	6.4 (109)	154	3.2 (102)	0	* *	0	* *	0	* *	0	* *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in

(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

(2)

- (1) Reporting period is January 1 through December 31.
- (2) TV Viewing and Smoking in Household data are not reported by all contributors.
- (3) Analyses based on one record per child.
- (4) Excludes records with unknown data or errors.
- (5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

- (6) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2006 Pediatric Nutrition Surveillance (1)
Michigan

Page: 17
Run Date: 5 / 15 / 07

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (4,5)							
	Breastfeeding (5)			Exclusive Breastfeeding (6)		≤2 Hours TV Viewing/Day (7)	Smoking in Household
Clinic	Number % Ever Breastfed (Rank)	Number % Breastfed At Least 6 Months (Rank)	Number % Breastfed At Least 12 Months (Rank)	Number % Breastfed At Least 3 Months (Rank)	Number % Breastfed At Least 6 Months (Rank)	Number % (Rank)	Number % (Rank)
Nation (Prior Year)	1,515,954 58.5	751,259 24.3	869,342 17.7	101,887 13.6	136,891 6.8	343,170 64.7	1,120,034 16.9

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in

(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2004 - 2006 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 1
Run Date: 5 / 15 / 07

Comparison of Growth and Anemia Indicators by Clinic (3,4)											
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)	
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number % (Rank)
0101	99	* *	* *	564	5.5 (61)	8.0(197)	11.5 (70)	254	10.6 (11)	9.8 (27)	382 8.9 (69)
0102	359	6.7 (36)	14.5 (171)	1,815	6.0 (85)	3.0 (75)	15.5 (173)	793	21.1 (179)	16.1 (153)	1,181 13.5 (126)
0103	414	9.7 (118)	6.5 (47)	1,935	6.4 (107)	3.3 (93)	14.2 (149)	846	18.8 (149)	12.8 (89)	1,260 9.8 (89)
0104	124	8.1 (77)	9.7 (123)	631	7.1 (138)	2.2 (27)	14.3 (150)	273	16.8 (101)	13.6 (115)	438 14.4 (132)
0250	174	6.3 (27)	10.3 (135)	830	3.9 (15)	2.0 (21)	16.5 (180)	394	13.2 (27)	19.0 (179)	478 14.0 (128)
0251	130	5.4 (15)	5.4 (25)	658	5.3 (50)	2.7 (58)	14.1 (145)	334	21.9 (183)	16.2 (156)	445 12.6 (117)
0252	137	6.6 (33)	8.0 (81)	656	5.2 (46)	2.4 (36)	17.5 (190)	315	24.8 (190)	18.7 (177)	370 15.1 (137)
0253	167	10.2 (129)	11.4 (156)	882	5.4 (56)	6.1 (183)	11.3 (68)	447	16.1 (74)	16.8 (163)	594 17.3 (153)
0401	526	11.2 (143)	11.0 (150)	2,142	7.9 (161)	1.4 (6)	16.6 (181)	943	17.7 (123)	14.1 (125)	1,346 8.1 (60)
0402	493	7.5 (63)	10.1 (131)	2,039	5.1 (40)	2.6 (49)	11.0 (52)	887	17.7 (123)	13.1 (99)	1,349 10.0 (92)
0403	154	7.1 (48)	13.0 (167)	764	3.0 (4)	6.4 (189)	10.2 (32)	392	10.7 (12)	10.7 (39)	516 16.3 (146)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2004 - 2006 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 2
Run Date: 5 / 15 / 07

Comparison of Growth and Anemia Indicators by Clinic (3,4)											
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)	
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number % (Rank)
0404	156	5.1 (13)	10.3 (135)	819	3.5 (10)	3.4 (96)	11.2 (62)	398	13.8 (32)	10.1 (34)	568 11.3 (105)
0501	298	4.7 (7)	12.1 (160)	2,213	3.3 (7)	6.4 (189)	6.9 (3)	1,164	9.9 (9)	5.6 (4)	1,663 4.1 (9)
0502	3	* *	* *	22	* *	* *	* *	9	* *	* *	15 * *
0503	150	7.3 (57)	8.7 (98)	687	5.4 (56)	5.1 (148)	14.7 (160)	305	13.8 (32)	11.8 (59)	461 4.6 (16)
0504	618	8.1 (77)	9.1 (108)	2,870	5.0 (34)	2.3 (33)	13.2 (117)	1,366	18.2 (137)	13.4 (111)	1,949 4.4 (11)
0505	446	7.4 (59)	10.5 (143)	1,927	6.0 (85)	3.7 (111)	12.6 (106)	930	16.7 (94)	14.0 (122)	1,341 5.5 (26)
0506	201	4.0 (3)	9.0 (105)	986	4.8 (28)	4.9 (146)	12.8 (113)	487	14.2 (41)	15.2 (143)	715 3.6 (5)
0650	1,036	7.9 (72)	8.5 (91)	4,069	7.7 (158)	3.1 (81)	10.6 (42)	1,666	19.9 (167)	13.0 (97)	2,428 6.7 (41)
0651	625	7.0 (46)	6.7 (53)	2,690	5.2 (46)	3.6 (106)	12.6 (106)	1,226	15.3 (61)	14.8 (131)	1,744 12.6 (117)
0652	411	5.8 (23)	10.0 (129)	1,775	6.1 (93)	2.6 (49)	13.5 (126)	819	16.1 (74)	17.1 (166)	1,200 10.6 (97)
0653	427	5.6 (20)	8.4 (86)	2,136	7.4 (152)	1.5 (9)	13.7 (132)	1,045	18.9 (151)	15.0 (139)	1,469 10.8 (100)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2004 - 2006 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 3
Run Date: 5 / 15 / 07

Comparison of Growth and Anemia Indicators by Clinic (3,4)													
Clinic	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)		
	Low		High	Short Stature (6)				Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
0654	351	6.3 (27)	15.7 (174)	1,593	4.4 (25)	3.8(115)	11.6 (74)	687	14.3 (43)	16.3(158)		1,041	9.0 (71)
0655	440	7.0 (46)	8.4 (86)	1,895	6.3 (99)	2.6 (49)	11.8 (83)	860	17.3(111)	11.6 (55)		1,277	6.3 (35)
0657	117	11.1 (142)	8.5 (91)	565	8.5(179)	4.1(126)	11.2 (62)	263	17.1(109)	13.3(107)		386	10.6 (97)
0701	393	3.1 (1)	10.9(149)	1,902	5.9 (78)	4.2(131)	8.3 (9)	861	12.5 (22)	6.5 (10)		1,132	9.4 (77)
0702	118	5.1 (13)	9.3(115)	497	5.0 (34)	5.8(177)	11.1 (58)	221	14.0 (38)	10.0 (31)		309	8.7 (65)
0703	254	6.3 (27)	11.0(150)	1,029	7.1(138)	2.9 (67)	9.7 (24)	449	18.0(131)	11.4 (50)		590	6.1 (29)
0704	89	* *	* *	382	7.3(150)	4.2(131)	10.7 (45)	167	10.8 (13)	6.0 (5)		245	4.9 (19)
0705	161	5.0 (10)	9.9(127)	1,020	5.5 (61)	2.5 (43)	12.5(104)	528	16.3 (85)	12.3 (74)		645	6.5 (39)
0850	1,012	9.4 (109)	8.6 (96)	4,265	7.7(158)	3.8(115)	11.7 (76)	1,835	20.3(171)	14.7(130)		2,666	3.8 (7)
0851	787	9.5(113)	10.8(147)	3,546	7.0(136)	5.1(148)	10.3 (35)	1,621	15.2 (58)	12.2 (70)		2,252	7.3 (47)
0901	1,930	7.7 (68)	10.4(140)	7,726	8.1(169)	2.7 (58)	11.2 (62)	3,394	17.8(129)	13.3(107)		4,576	8.3 (62)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2004 - 2006 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)											
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)	
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number % (Rank)
0902	84	* *	* *	391	7.9(161)	2.0 (21)	11.8 (83)	184	25.5(191)	13.6(115)	257 10.9(101)
1051	1,425	9.0 (96)	10.2 (132)	5,868	6.3 (99)	5.1(148)	11.8 (83)	2,687	14.3 (43)	10.9 (44)	3,569 11.7 (109)
1059	1	* *	* *	35	* *	* *	* *	27	* *	* *	28 * *
1060	0	* *	* *	16	* *	* *	* *	13	* *	* *	14 * *
1101	1,504	9.2 (101)	6.4 (41)	7,230	6.3 (99)	3.1 (81)	11.2 (62)	3,193	16.0 (71)	12.1 (67)	4,960 17.6 (156)
1102	973	7.1 (48)	7.6 (69)	3,613	5.8 (75)	1.8 (17)	12.6(106)	1,478	15.3 (61)	11.5 (51)	2,249 11.9 (113)
1106	122	3.3 (2)	9.0 (105)	472	5.9 (78)	1.1 (5)	10.4 (39)	192	19.3(160)	6.8 (12)	312 16.3 (146)
1250	1,023	6.0 (25)	9.1 (108)	4,226	6.8(132)	5.2(155)	11.3 (68)	1,899	14.7 (48)	9.8 (27)	2,646 11.8 (111)
1251	482	5.4 (15)	10.4 (140)	3,076	5.7 (71)	4.5(141)	11.0 (52)	1,492	14.0 (38)	13.5(114)	2,205 10.9 (101)
1252	769	6.5 (32)	6.9 (56)	3,535	6.2 (98)	4.3(135)	19.3(196)	1,558	21.6(181)	28.8(191)	2,333 15.5 (140)
1253	796	8.7 (88)	6.8 (55)	3,206	6.7(124)	4.1(126)	12.5(104)	1,353	17.3(111)	12.7 (82)	2,000 9.4 (77)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2004 - 2006 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 5
Run Date: 5 / 15 / 07

Comparison of Growth and Anemia Indicators by Clinic (3,4)											
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)	
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number % (Rank)
1350	2,533	8.6 (87)	9.1 (108)	10,048	8.6(181)	3.5(100)	13.2(117)	4,205	19.1(155)	14.9(134)	6,456 18.1 (159)
1351	182	5.5 (18)	13.2 (168)	872	8.1(169)	1.6 (12)	14.1(145)	386	17.4(116)	13.2(104)	581 9.3 (75)
1352	483	8.5 (85)	7.5 (68)	2,120	8.4(178)	4.4(136)	12.3 (98)	955	18.7(146)	13.1 (99)	1,374 11.7 (109)
1401	263	10.6 (137)	9.1 (108)	1,177	5.9 (78)	5.6(171)	12.9(114)	513	17.3(111)	12.7 (82)	714 6.7 (41)
1402	174	6.9 (42)	15.5 (173)	797	5.3 (50)	2.9 (67)	19.9(198)	361	18.0(131)	23.3(183)	538 10.0 (92)
1403	23	* *	* *	187	13.4(207)	10.7(206)	17.6(191)	115	14.8 (50)	11.3 (47)	124 2.4 (3)
1404	12	* *	* *	90	* *	* *	* *	44	* *	* *	77 * *
1405	0	* *	* *	2	* *	* *	* *	2	* *	* *	2 * *
1501	1,111	6.8 (38)	8.8 (100)	5,161	5.9 (78)	4.0(121)	16.3(179)	2,423	18.9(151)	16.9(165)	3,327 15.2 (138)
1502	101	11.9 (146)	8.9 (103)	412	4.4 (25)	5.6(171)	12.4(102)	179	8.9 (6)	16.8(163)	246 13.4 (124)
1503	1,148	10.0 (125)	8.8 (100)	5,120	6.8(132)	4.5(141)	13.5(126)	2,364	15.1 (55)	14.8(131)	3,417 16.1 (145)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2004 - 2006 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 6
Run Date: 5 / 15 / 07

Comparison of Growth and Anemia Indicators by Clinic (3,4)													
Clinic	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)		Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
1504	2,214	10.1 (127)	7.6 (69)	10,068	9.2(192)	2.8 (62)	13.3(121)	4,441	14.9 (52)	12.6 (80)		6,424	6.5 (39)
1505	625	9.9 (123)	11.2 (154)	2,837	5.7 (71)	4.8(145)	9.8 (26)	1,308	13.5 (30)	8.2 (16)		1,846	8.7 (65)
1506	498	6.8 (38)	12.2 (162)	2,076	7.9(161)	5.1 (148)	15.7 (176)	918	18.4 (143)	13.2 (104)		1,307	18.7 (162)
1507	565	7.4 (59)	12.4 (166)	2,887	6.4(107)	4.0(121)	13.7 (132)	1,390	18.1 (134)	14.1 (125)		1,880	9.5 (83)
1508	155	7.1 (48)	14.2 (170)	904	14.0(208)	5.4(163)	20.9(199)	449	20.3(171)	23.8(184)		573	18.7 (162)
1509	516	10.9 (140)	7.6 (69)	2,999	7.9(161)	2.5 (43)	15.2(169)	1,415	16.6 (93)	12.7 (82)		2,021	22.3 (175)
1510	171	9.4 (109)	6.4 (41)	996	6.6(117)	3.0 (75)	11.9 (88)	429	17.7(123)	9.1 (21)		675	17.3 (153)
1511	559	5.7 (21)	10.7 (146)	2,167	6.4(107)	5.7(175)	12.3 (98)	897	18.8(149)	14.9(134)		1,337	22.1 (173)
1512	318	11.3 (145)	7.9 (78)	1,221	10.6(203)	3.1 (81)	16.0(177)	543	18.2(137)	14.9(134)		744	20.8 (169)
1701	698	6.7 (36)	13.8 (169)	2,790	6.0 (85)	2.3 (33)	15.6(175)	1,227	19.3(160)	17.2(167)		1,847	9.4 (77)
1712	129	4.7 (7)	9.3 (115)	689	5.1 (40)	2.2 (27)	13.6(130)	362	17.4(116)	12.7 (82)		499	14.0 (128)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2004 - 2006 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 7
Run Date: 5 / 15 / 07

Comparison of Growth and Anemia Indicators by Clinic (3,4)												
Clinic	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)	
	Low		High	Short Stature (6)		Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
1950	701	8.3 (82)	7.0 (59)	2,968	5.9 (78)	5.2(155)	11.2 (62)	1,325	9.7 (8)	8.2 (16)	1,869	6.2 (32)
1952	175	10.3 (131)	9.7 (123)	750	6.7(124)	2.5 (43)	14.4(153)	337	17.5(119)	15.1(141)	481	7.5 (50)
1953	389	8.7 (88)	8.7 (98)	1,502	5.7 (71)	5.1(148)	8.7 (11)	618	11.5 (16)	6.5 (10)	917	6.8 (44)
1954	377	9.0 (96)	8.5 (91)	1,694	8.2(174)	2.5 (43)	16.6(181)	753	17.3(111)	17.9(173)	1,040	5.2 (24)
1955	191	4.2 (5)	8.4 (86)	856	8.6(181)	3.0 (75)	15.1(167)	417	15.8 (69)	14.6(127)	527	6.1 (29)
1956	14	* *	* *	55	* *	* *	* *	25	* *	* *	28	* *
1957	377	7.7 (68)	9.5 (121)	1,720	4.9 (31)	4.1(126)	10.8 (48)	740	7.2 (4)	6.2 (8)	1,072	4.6 (16)
1959	196	9.2 (101)	6.6 (49)	892	7.1(138)	3.0 (75)	10.7 (45)	379	14.0 (38)	9.2 (23)	544	4.8 (18)
1963	205	8.3 (82)	5.9 (30)	831	8.8(186)	1.4 (6)	15.0(164)	334	19.2(158)	15.6(147)	478	5.0 (21)
2150	739	8.3 (82)	11.4 (156)	3,323	5.4 (56)	2.4 (36)	12.2 (95)	1,587	18.3(142)	16.4(160)	2,162	2.1 (1)
2151	327	5.5 (18)	6.4 (41)	1,433	5.5 (61)	2.4 (36)	8.9 (13)	669	16.3 (85)	12.1 (67)	908	2.2 (2)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2004 - 2006 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 8
Run Date: 5 / 15 / 07

Comparison of Growth and Anemia Indicators by Clinic (3,4)												
Clinic	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)	
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
2251	549	6.6 (33)	10.2 (132)	2,365	5.6 (67)	1.7 (15)	13.2(117)	1,087	18.9(151)	15.4(144)	1,526	7.8 (56)
2252	197	4.1 (4)	12.2 (162)	911	6.1 (93)	4.6(143)	13.4(123)	418	17.9(130)	17.5(171)	592	10.0 (92)
2551	3,963	12.6 (154)	4.3 (5)	15,238	6.1 (93)	4.4(136)	10.1 (29)	6,226	15.4 (64)	12.7 (82)	9,410	24.6 (182)
2552	880	12.5 (152)	5.9 (30)	3,707	4.9 (31)	4.6(143)	10.2 (32)	1,482	16.7 (94)	13.3(107)	2,239	28.1 (188)
2554	100	5.0 (10)	12.0 (159)	607	7.1 (138)	5.8(177)	11.7 (76)	320	15.0 (53)	9.4 (26)	375	14.4 (132)
2555	3,904	9.8 (122)	7.7 (74)	15,563	6.0 (85)	4.4(136)	11.5 (70)	6,213	15.2 (58)	12.9 (93)	9,351	19.2 (166)
3001	1,105	14.3 (160)	5.7 (26)	4,974	6.4 (107)	2.9 (67)	15.2(169)	2,543	16.7 (94)	12.2 (70)	3,410	13.3 (122)
3002	222	12.2 (149)	4.5 (7)	1,062	4.3 (20)	2.9 (67)	16.7(183)	566	16.3 (85)	12.9 (93)	764	29.1 (190)
3003	21	* *	* *	127	7.9(161)	0.8 (3)	15.0(164)	81	* *	* *	96	* *
3004	244	6.6 (33)	5.7 (26)	1,288	5.0 (34)	1.6 (12)	24.0(204)	697	16.4 (89)	26.1(188)	859	7.3 (47)
3007	34	* *	* *	177	2.3 (3)	2.8 (62)	18.6(195)	91	* *	* *	135	8.1 (60)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2004 - 2006 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)													
Clinic	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)		
	Low		High	Short Stature (6)		Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
3009	5	* *	* *	49	* *	* *	* *	31	* *	* *		44	* *
3010	1	* *	* *	13	* *	* *	* *	9	* *	* *		11	* *
3201	604	8.1 (77)	9.8 (126)	2,690	5.3 (50)	4.0(121)	12.3 (98)	1,227	16.1 (74)	13.9(121)		1,703	9.7 (88)
3202	13	* *	* *	100	2.0 (2)	0.0 (1)	15.0(164)	61	* *	* *		66	* *
3301	2,280	9.6 (115)	8.6 (96)	14,106	6.3 (99)	2.9 (67)	13.1(116)	6,313	17.0(105)	13.0 (97)		9,921	13.4 (124)
3302	259	10.4 (132)	6.2 (37)	2,050	5.6 (67)	3.5(100)	12.2 (95)	1,069	17.0(105)	13.7(118)		1,531	13.5 (126)
3303	47	* *	* *	290	6.6(117)	3.1 (81)	9.0 (15)	130	17.7(123)	10.8 (41)		207	7.7 (53)
3304	88	* *	* *	682	3.4 (8)	3.5(100)	7.5 (5)	317	12.0 (20)	6.0 (5)		504	11.3 (105)
3305	57	* *	* *	500	5.6 (67)	2.6 (49)	9.8 (26)	232	21.6(181)	12.1 (67)		369	9.5 (83)
3306	63	* *	* *	353	4.0 (17)	4.2(131)	9.9 (28)	127	20.5(174)	15.7(148)		260	6.2 (32)
3308	38	* *	* *	235	6.0 (85)	3.0 (75)	11.9 (88)	108	18.5(144)	17.6(172)		166	7.8 (56)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2004 - 2006 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)												
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
3309	25	* *	* *	164	4.9 (31)	3.7(111)	9.1 (16)	72	* *	* *	111	6.3 (35)
3311	12	* *	* *	34	* *	* *	* *	5	* *	* *	19	* *
3550	98	* *	* *	391	5.4 (56)	2.8 (62)	17.1(187)	173	22.0(184)	13.3(107)	232	18.5 (160)
3551	35	* *	* *	138	7.2(145)	2.2 (27)	16.7(183)	54	* *	* *	71	* *
3552	40	* *	* *	217	6.5(113)	3.7(111)	25.3(207)	108	15.7 (67)	25.0(186)	137	26.3 (185)
3650	24	* *	* *	112	7.1(138)	0.0 (1)	25.0(206)	54	* *	* *	87	* *
3701	774	7.2 (52)	10.6(144)	2,959	6.6(117)	5.4(163)	13.5(126)	1,249	21.0(177)	18.7(177)	1,908	7.9 (59)
3702	251	12.0(147)	8.0 (81)	881	6.7(124)	4.0(121)	14.0(143)	353	17.6(121)	18.1(174)	542	9.4 (77)
3703	126	12.7(155)	4.8 (12)	406	5.9 (78)	6.2(185)	14.5(157)	135	20.7(176)	27.4(190)	237	11.0(103)
3704	105	5.7 (21)	11.4(156)	383	8.1(169)	3.4 (96)	11.0 (52)	145	21.4(180)	15.9(150)	243	5.3 (25)
3805	3,372	9.2(101)	7.4 (64)	12,311	6.7(124)	3.2 (89)	10.9 (50)	4,893	16.7 (94)	14.0(122)	6,296	7.8 (56)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2004 - 2006 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)												
Clinic	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)	
	Low		High	Short Stature (6)			Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
3950	1,905	10.7 (139)	10.3 (135)	8,405	7.2(145)	5.4(163)	14.1(145)	3,570	18.0(131)	15.4(144)	5,163	23.0 (178)
3952	64	* *	* *	327	8.6(181)	3.1 (81)	12.2 (95)	141	14.2 (41)	9.2 (23)	214	15.0 (136)
3953	90	* *	* *	364	6.3 (99)	7.1(193)	11.0 (52)	157	15.3 (61)	14.0(122)	228	17.1 (151)
3954	173	6.9 (42)	11.0 (150)	812	5.5 (61)	5.5(166)	14.3(150)	351	19.7(164)	18.2(176)	503	17.5 (155)
4001	999	12.9 (156)	4.8 (12)	4,347	6.0 (85)	5.5(166)	12.6(106)	1,959	18.2(137)	13.4(111)	2,983	24.3 (181)
4002	799	10.6 (137)	7.6 (69)	3,425	7.6(156)	6.2(185)	11.5 (70)	1,491	16.2 (81)	11.8 (59)	2,294	20.7 (168)
4103	39	* *	* *	296	3.7 (12)	8.8(201)	9.1 (16)	173	15.6 (66)	9.2 (23)	225	24.0 (180)
4106	1,618	10.4 (132)	5.1 (18)	7,505	6.5(113)	2.1 (24)	13.8(138)	3,607	17.5(119)	15.9(150)	5,152	10.1 (95)
4107	2,631	12.0 (147)	8.0 (81)	12,565	8.2(174)	2.7 (58)	13.3(121)	5,741	16.8(101)	11.9 (62)	8,268	6.0 (28)
4109	1,127	9.3 (107)	10.0 (129)	4,225	6.5(113)	2.5 (43)	10.1 (29)	1,805	16.1 (74)	11.3 (47)	2,670	2.5 (4)
4110	1,621	7.6 (66)	9.3 (115)	7,059	6.6(117)	1.5 (9)	14.5(157)	3,293	20.6(175)	17.4(169)	4,705	4.4 (11)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2004 - 2006 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 12
Run Date: 5 / 15 / 07

Comparison of Growth and Anemia Indicators by Clinic (3,4)												
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low		High	Short Stature (6)		Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
4111	63	* *	* *	200	9.5(196)	3.5(100)	14.0(143)	75	* *	* *	114	12.3 (114)
4112	808	4.7 (7)	10.3 (135)	3,271	8.5(179)	2.4 (36)	21.5(202)	1,486	19.1 (155)	26.7(189)	2,296	9.1 (74)
4113	495	4.6 (6)	9.7 (123)	2,920	4.2 (18)	3.4 (96)	14.8(161)	1,453	16.2 (81)	20.6(180)	2,112	11.4 (108)
4120	1,971	7.9 (72)	9.6 (122)	8,049	6.6(117)	2.5 (43)	12.4(102)	3,416	16.0 (71)	12.2 (70)	5,100	6.2 (32)
4160	1,864	8.9 (94)	9.2 (113)	7,410	6.7(124)	3.2 (89)	13.4(123)	3,133	16.1 (74)	13.6(115)	4,530	4.4 (11)
4351	733	7.5 (63)	7.8 (76)	3,031	6.9(135)	2.4 (36)	13.7(132)	1,322	16.4 (89)	12.0 (65)	1,930	9.8 (89)
4352	223	7.6 (66)	9.9 (127)	907	7.1(138)	3.6(106)	11.2 (62)	394	19.0(154)	11.7 (57)	589	21.7 (172)
4353	531	8.5 (85)	7.9 (78)	2,288	8.7(185)	1.8 (17)	13.2(117)	999	20.1(169)	14.6(127)	1,494	8.6 (64)
4354	818	10.4 (132)	12.1 (160)	3,353	4.7 (27)	4.4(136)	8.1 (8)	1,434	12.6 (24)	9.9 (30)	2,105	12.5 (116)
4355	381	9.7 (118)	6.3 (39)	1,572	5.3 (50)	5.5(166)	8.0 (6)	703	11.5 (16)	9.0 (20)	1,014	6.8 (44)
4356	588	8.2 (80)	7.7 (74)	2,384	4.8 (28)	5.5(166)	7.4 (4)	1,051	9.9 (9)	8.2 (16)	1,547	9.0 (71)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2004 - 2006 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)											
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)	
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number % (Rank)
4357	13	* *	* *	62	* *	* *	* *	32	* *	* *	39 * *
4358	239	5.4 (15)	8.4 (86)	1,324	7.2(145)	1.5 (9)	21.2(201)	677	23.0(186)	24.7(185)	968 9.9 (91)
4359	360	7.2 (52)	10.6(144)	1,504	10.0(200)	2.1 (24)	18.3(193)	719	18.2(137)	17.4(169)	960 15.8(143)
4360	212	6.1 (26)	10.4(140)	797	4.3 (20)	2.8 (62)	8.8 (12)	362	11.0 (15)	8.8 (19)	503 13.1(120)
4361	271	7.7 (68)	6.3 (39)	1,101	5.7 (71)	1.6 (12)	10.3 (35)	495	20.0(168)	12.9 (93)	716 7.7 (53)
4362	396	6.8 (38)	6.6 (49)	1,618	7.2(145)	3.0 (75)	10.6 (42)	703	16.9(104)	11.9 (62)	1,021 10.6 (97)
4363	879	7.2 (52)	10.8(147)	3,516	5.0 (34)	2.2 (27)	9.5 (21)	1,472	18.1(134)	11.2 (46)	2,202 11.1(104)
4401	1,209	7.9 (72)	9.3(115)	4,791	9.6(198)	3.1 (81)	12.9(114)	2,066	16.7 (94)	12.9 (93)	3,167 9.3 (75)
4501	1,067	8.7 (88)	9.2(113)	4,796	6.3 (99)	3.1 (81)	11.0 (52)	2,159	15.1 (55)	11.5 (51)	3,183 5.0 (21)
4601	1,523	8.7 (88)	6.1 (36)	5,803	5.2 (46)	4.2(131)	13.8(138)	2,431	16.2 (81)	15.1(141)	2,851 14.0(128)
4701	1,067	8.2 (80)	9.3(115)	3,833	5.3 (50)	3.2 (89)	10.7 (45)	1,434	16.4 (89)	11.9 (62)	2,193 5.5 (26)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2004 - 2006 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 14
Run Date: 5 / 15 / 07

Comparison of Growth and Anemia Indicators by Clinic (3,4)													
Clinic	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)		Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
5050	1,522	13.1 (157)	6.4 (41)	4,872	7.5(154)	2.2 (27)	14.5(157)	1,465	14.7 (48)	12.6 (80)		2,551	21.4 (171)
5051	3,856	10.2 (129)	6.9 (56)	12,902	6.8(132)	2.4 (36)	14.9(163)	4,056	12.8 (25)	10.6 (38)		7,264	16.7 (150)
5052	2,311	9.4 (109)	6.4 (41)	7,801	6.7(124)	2.0 (21)	14.4(153)	2,493	13.1 (26)	13.7(118)		4,366	14.7 (134)
5054	251	7.2 (52)	5.2 (21)	936	8.1(169)	1.4 (6)	23.3(203)	303	13.9 (36)	21.8(182)		525	11.8 (111)
5202	332	7.2 (52)	10.2 (132)	1,459	8.0(167)	2.2 (27)	11.7 (76)	635	19.8(166)	12.0 (65)		817	6.7 (41)
5203	188	7.4 (59)	9.0 (105)	925	8.9(187)	2.7 (58)	13.5(126)	463	21.0(177)	14.9(134)		561	4.5 (15)
5204	159	6.9 (42)	14.5 (171)	617	7.3(150)	2.1 (24)	12.0 (92)	258	23.6(187)	10.1 (34)		349	8.3 (62)
5205	255	12.5 (152)	8.2 (84)	1,050	9.4(194)	1.8 (17)	13.8(138)	479	18.6(145)	16.1 (153)		613	4.2 (10)
5601	1,136	9.6 (115)	9.1 (108)	4,780	9.1(190)	1.8 (17)	21.0(200)	2,167	18.7(146)	16.1 (153)		2,986	9.6 (87)
5801	2,068	9.3 (107)	6.0 (32)	6,855	7.5(154)	2.6 (49)	11.1 (58)	2,328	14.5 (47)	12.5 (79)		3,605	9.5 (83)
5803	16	* *	* *	120	5.0 (34)	0.8 (3)	16.7(183)	85	* *	* *		73	* *

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2004 - 2006 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 15
Run Date: 5 / 15 / 07

Comparison of Growth and Anemia Indicators by Clinic (3,4)												
Clinic	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)	
	Low		High	Short Stature (6)		Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
6103	4,376	9.0 (96)	7.4 (64)	17,695	6.0 (85)	5.6(171)	10.5 (40)	8,136	18.2(137)	12.8 (89)	11,648	16.3 (146)
6301	4,781	9.6 (115)	7.3 (62)	18,084	6.7(124)	2.9 (67)	11.1 (58)	7,453	16.1 (74)	13.8(120)	11,453	14.8 (135)
6312	2,748	9.9 (123)	7.9 (78)	10,629	5.4 (56)	5.7(175)	9.2 (18)	4,329	10.9 (14)	7.2 (14)	6,913	17.1 (151)
6316	1,170	10.0 (125)	7.6 (69)	4,188	5.0 (34)	3.3 (93)	8.0 (6)	1,578	17.0(105)	9.1 (21)	2,609	9.0 (71)
6320	252	6.3 (27)	10.3 (135)	1,109	5.1 (40)	7.3(194)	9.6 (22)	469	15.1 (55)	11.7 (57)	682	8.7 (65)
6325	141	5.0 (10)	11.3 (155)	718	6.0 (85)	3.6(106)	11.0 (52)	343	17.2(110)	13.1 (99)	454	8.8 (68)
6330	250	13.2 (158)	6.0 (32)	967	11.1(205)	5.6(171)	11.7 (76)	385	14.3 (43)	12.7 (82)	578	12.8 (119)
6335	582	8.9 (94)	7.0 (59)	2,530	5.1 (40)	5.5(166)	10.3 (35)	983	14.8 (50)	13.1 (99)	1,543	9.5 (83)
6340	484	7.4 (59)	6.4 (41)	1,750	6.7(124)	4.1(126)	10.9 (50)	569	16.0 (71)	12.7 (82)	974	10.4 (96)
7304	312	6.4 (31)	12.2 (162)	1,234	5.1 (40)	2.4 (36)	15.2(169)	599	22.0(184)	16.7(161)	909	6.1 (29)
7306	3,019	10.5 (136)	8.5 (91)	11,300	5.5 (61)	2.6 (49)	13.4(123)	5,443	16.2 (81)	12.4 (77)	8,217	9.4 (77)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2004 - 2006 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 16
Run Date: 5 / 15 / 07

Comparison of Growth and Anemia Indicators by Clinic (3,4)											
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)	
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number % (Rank)
7308	227	7.9 (72)	12.3 (165)	843	5.1 (40)	2.8 (62)	14.1 (145)	425	19.1 (155)	16.2 (156)	648 4.9 (19)
7401	2,465	8.8 (93)	6.7 (53)	9,458	5.9 (78)	8.4 (199)	10.6 (42)	4,036	14.4 (46)	11.3 (47)	6,136 12.4 (115)
7402	352	9.1 (99)	8.2 (84)	1,392	5.8 (75)	3.5 (100)	10.3 (35)	577	20.1 (169)	11.6 (55)	887 6.8 (44)
7403	99	* *	* *	375	6.4 (107)	5.3 (160)	13.6 (130)	143	23.8 (188)	15.4 (144)	258 8.9 (69)
7404	145	6.9 (42)	3.4 (2)	545	5.3 (50)	8.8 (201)	11.9 (88)	214	18.7 (146)	10.7 (39)	345 11.3 (105)
7601	776	7.7 (68)	7.2 (61)	3,416	3.8 (13)	5.3 (160)	8.3 (9)	1,568	17.3 (111)	11.5 (51)	2,224 5.0 (21)
7850	1,148	7.5 (63)	7.4 (64)	4,351	6.1 (93)	7.3 (194)	9.6 (22)	1,906	15.8 (69)	12.2 (70)	2,774 6.3 (35)
7851	220	5.9 (24)	7.3 (62)	1,086	6.3 (99)	5.2 (155)	10.1 (29)	517	17.6 (121)	11.8 (59)	747 3.6 (5)
7852	126	7.1 (48)	11.1 (153)	566	5.5 (61)	3.9 (119)	11.8 (83)	276	17.0 (105)	16.3 (158)	384 4.4 (11)
8150	83	* *	* *	393	5.6 (67)	5.1 (148)	11.7 (76)	166	18.1 (134)	15.7 (148)	214 15.4 (139)
8151	2,524	9.2 (101)	8.8 (100)	9,730	6.6 (117)	5.1 (148)	12.7 (111)	4,029	17.7 (123)	14.9 (134)	6,062 26.8 (186)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2004 - 2006 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 17
Run Date: 5 / 15 / 07

Comparison of Growth and Anemia Indicators by Clinic (3,4)											
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)	
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number % (Rank)
8152	495	9.7 (118)	4.0 (3)	1,358	3.5 (10)	8.5(200)	10.2 (32)	333	19.5(163)	10.8 (41)	649 22.2 (174)
8153	36	* *	* *	131	6.1 (93)	3.8(115)	16.8(186)	46	* *	* *	73 * *
8154	364	8.0 (76)	5.2 (21)	1,560	3.8 (13)	6.2(185)	6.7 (1)	694	12.2 (21)	6.3 (9)	915 9.4 (77)
8155	71	* *	* *	377	5.8 (75)	5.8(177)	9.3 (19)	204	11.8 (19)	9.8 (27)	268 17.9 (158)
8156	38	* *	* *	175	8.0(167)	6.3(188)	15.4(172)	72	* *	* *	95 * *
8157	5	* *	* *	59	* *	* *	* *	40	* *	* *	40 * *
8201	2,557	9.2 (101)	6.9 (56)	8,021	6.4(107)	3.3 (93)	11.7 (76)	2,695	15.2 (58)	12.3 (74)	4,365 7.5 (50)
8202	2,626	7.3 (57)	7.4 (64)	12,499	4.2 (18)	4.1(126)	11.5 (70)	5,621	15.0 (53)	13.1 (99)	7,789 7.7 (53)
8253	417	9.1 (99)	9.4 (120)	1,753	7.2(145)	2.6 (49)	14.3(150)	649	20.3(171)	12.8 (89)	1,009 16.0 (144)
8254	483	8.7 (88)	8.9 (103)	2,150	5.2 (46)	3.1 (81)	13.7(132)	864	17.7(123)	14.8(131)	1,194 3.9 (8)
8255	3,058	9.4 (109)	8.5 (91)	9,913	8.6(181)	3.6(106)	11.9 (88)	3,493	17.4(116)	14.6(127)	5,167 7.4 (49)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2004 - 2006 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 18
Run Date: 5 / 15 / 07

Comparison of Growth and Anemia Indicators by Clinic (3,4)											
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)	
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number % (Rank)
8257	481	11.2 (143)	6.0 (32)	1,453	9.5(196)	5.2(155)	15.1(167)	585	13.8 (32)	13.2(104)	929 27.0 (187)
8258	1,792	9.7 (118)	7.8 (76)	6,452	7.1(138)	2.3 (33)	14.8(161)	2,655	19.7(164)	16.0(152)	4,069 17.6 (156)
8901	1,002	14.9 (163)	4.0 (3)	4,682	9.6(198)	5.9(180)	13.7(132)	2,200	16.1 (74)	11.5 (51)	1,204 22.4 (176)
8902	516	12.4 (151)	5.0 (17)	2,280	7.4(152)	6.5(192)	11.8 (83)	1,041	13.5 (30)	10.5 (37)	407 18.9 (164)
8903	1,130	15.8 (169)	3.0 (1)	4,669	9.0(188)	5.3(160)	14.4(153)	2,039	16.3 (85)	10.3 (36)	1,657 25.5 (183)
8904	694	12.2 (149)	5.8 (28)	3,767	6.5(113)	8.3(198)	13.9(141)	1,831	13.3 (28)	13.4(111)	2,271 23.5 (179)
8905	1,899	10.1 (127)	6.6 (49)	7,295	7.7(158)	7.9(196)	16.2(178)	3,159	16.8(101)	16.7(161)	529 6.4 (38)
9001	3	* *	* *	56	* *	* *	* *	27	* *	* *	42 * *
9002	7	* *	* *	121	8.3(177)	1.7 (15)	28.1(208)	63	* *	* *	95 * *
9004	193	18.7 (174)	4.7 (10)	1,607	10.5(202)	2.6 (49)	24.6(205)	851	24.1 (189)	25.4(187)	1,227 32.5 (192)
9005	4	* *	* *	42	* *	* *	* *	27	* *	* *	32 * *

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2004 - 2006 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 19
Run Date: 5 / 15 / 07

Comparison of Growth and Anemia Indicators by Clinic (3,4)											
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)	
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number % (Rank)
9006	9	* *	* *	135	3.0 (4)	12.6(207)	12.6(106)	79	* *	* *	117 47.9 (197)
9008	2	* *	* *	41	* *	* *	* *	24	* *	* *	31 * *
9009	15	* *	* *	124	4.8 (28)	4.0(121)	10.5 (40)	65	* *	* *	86 * *
9010	179	18.4 (173)	4.5 (7)	1,117	7.0(136)	4.4(136)	15.5(173)	597	16.4 (89)	12.4 (77)	722 36.1 (194)
9011	4	* *	* *	48	* *	* *	* *	29	* *	* *	37 * *
9012	7	* *	* *	116	4.3 (20)	2.6 (49)	12.1 (93)	81	* *	* *	91 * *
9014	6	* *	* *	66	* *	* *	* *	37	* *	* *	57 * *
9018	0	* *	* *	35	* *	* *	* *	16	* *	* *	28 * *
9020	6	* *	* *	117	0.9 (1)	3.4 (96)	17.9(192)	77	* *	* *	94 * *
9022	4	* *	* *	43	* *	* *	* *	22	* *	* *	30 * *
9023	10	* *	* *	97	* *	* *	* *	58	* *	* *	58 * *

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2004 - 2006 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)											
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)	
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number % (Rank)
9024	5	* *	* *	83	* *	* *	* *	48	* *	* *	71 * *
9035	9	* *	* *	161	4.3 (20)	3.7(111)	13.7(132)	96	* *	* *	127 37.8 (195)
9039	12	* *	* *	152	3.9 (15)	5.9(180)	17.1(187)	83	* *	* *	118 33.1 (193)
9040	133	15.8 (169)	6.0 (32)	1,090	3.2 (6)	3.8(115)	10.8 (48)	503	12.5 (22)	7.0 (13)	809 29.2 (191)
9045	1,250	10.4 (132)	6.2 (37)	5,329	10.4(201)	6.1(183)	19.4(197)	2,424	19.2(158)	20.9(181)	842 7.6 (52)
9050	1,859	9.5 (113)	6.6 (49)	7,881	6.3 (99)	5.9(180)	12.3 (98)	3,447	11.7 (18)	10.0 (31)	4,628 16.4 (149)
9051	2,070	6.8 (38)	8.4 (86)	9,330	6.6(117)	2.9 (67)	17.1(187)	4,064	15.4 (64)	17.3(168)	6,051 15.7 (142)
9053	2,498	14.3 (160)	4.9 (15)	9,456	9.3(193)	6.4(189)	11.1 (58)	3,667	9.4 (7)	7.4 (15)	6,080 18.6 (161)
9054	640	15.8 (169)	5.2 (21)	1,960	7.6(156)	5.0(147)	12.7(111)	641	16.7 (94)	12.8 (89)	1,124 42.0 (196)
9058	487	9.2 (101)	5.3 (24)	2,311	3.4 (8)	19.1(208)	6.8 (2)	1,093	5.3 (1)	3.7 (1)	1,683 20.4 (167)
9066	2,397	15.9 (172)	4.4 (6)	8,734	10.7(204)	9.0(203)	8.9 (13)	3,358	7.9 (5)	6.0 (5)	4,030 14.3 (131)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2004 - 2006 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 21
Run Date: 5 / 15 / 07

Comparison of Growth and Anemia Indicators by Clinic (3,4)												
Clinic	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)	
	Low		High	Short Stature (6)		Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
9068	1,425	15.6 (167)	5.8 (28)	5,748	8.2(174)	9.5(205)	9.7 (24)	2,463	6.7 (3)	4.3 (2)	3,333	22.5 (177)
9069	1,903	15.2 (164)	5.1 (18)	6,814	8.1(169)	3.9(119)	11.6 (74)	2,291	19.3(160)	15.0(139)	3,793	15.6 (141)
9070	1,730	14.5 (162)	4.9 (15)	7,541	9.0(188)	9.3(204)	13.9(141)	2,991	6.0 (2)	5.5 (3)	4,444	13.2 (121)
9072	1,599	15.6 (167)	4.7 (10)	5,695	9.4(194)	3.6(106)	11.7 (76)	2,218	13.8 (32)	10.8 (41)	2,791	13.3 (122)
9074	1,363	13.6 (159)	4.8 (12)	4,859	7.9(161)	3.2 (89)	12.1 (93)	1,932	13.9 (36)	11.0 (45)	3,016	19.1 (165)
9079	1,243	15.3 (166)	4.5 (7)	4,472	9.1(190)	3.5(100)	14.4(153)	1,680	13.3 (28)	12.3 (74)	2,439	26.1 (184)
9080	1,958	10.9 (140)	6.5 (47)	9,818	4.3 (20)	2.9 (67)	9.4 (20)	4,873	15.7 (67)	10.0 (31)	6,439	21.3 (170)
9150	0	* *	* *	10	* *	* *	* *	6	* *	* *	6	* *
9151	9	* *	* *	28	* *	* *	* *	10	* *	* *	15	* *
Unknown Clinics	257	15.2 (164)	5.1 (18)	1,248	12.3(206)	5.2(155)	18.4(194)	574	16.7 (94)	18.1(174)	723	28.1 (188)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2004 - 2006 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 22
Run Date: 5 / 15 / 07

Comparison of Growth and Anemia Indicators by Clinic (3,4)												
Clinic	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)	
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% >95th (Rank)	Number	% (Rank)
Michigan	161,795	9.8	7.7	669,540	6.8	4.1	12.5	286,870	15.9	13.0	408,408	13.3

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2004 - 2006 Pediatric Nutrition Surveillance (1)
Michigan

Page: 1
Run Date: 5 / 15 / 07

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
(4,5)

	Breastfeeding (5)						Exclusive Breastfeeding (6)				≤2 Hours TV Viewing/Day (7)	Smoking in Household		
Clinic	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months	Number	% Breastfed At Least 3 Months	Number	% Breastfed At Least 6 Months	Number	%	Number	%
	(Rank)		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)	
0101	151	51.7(112)	114	14.0 (98)	95	* *	0	* *	0	* *	0	* *	0	* *
0102	495	41.4(158)	396	7.6 (172)	263	6.5 (164)	0	* *	0	* *	0	* *	0	* *
0103	513	42.9(154)	403	6.5 (177)	312	5.8 (166)	0	* *	0	* *	0	* *	0	* *
0104	164	44.5(144)	127	10.2 (154)	104	8.7 (136)	0	* *	0	* *	0	* *	0	* *
0250	194	48.5(124)	175	11.4 (136)	158	7.6 (154)	0	* *	0	* *	0	* *	0	* *
0251	149	44.3(145)	130	10.8 (146)	104	12.5 (82)	0	* *	0	* *	0	* *	0	* *
0252	155	60.6 (51)	143	17.5 (58)	123	13.0 (71)	0	* *	0	* *	0	* *	0	* *
0253	193	59.6 (59)	168	25.6 (15)	150	25.3 (9)	0	* *	0	* *	0	* *	0	* *
0401	590	57.6 (75)	487	18.3 (49)	390	16.2 (41)	0	* *	0	* *	0	* *	0	* *
0402	551	65.9 (21)	452	16.6 (67)	380	13.9 (61)	0	* *	0	* *	0	* *	0	* *
0403	174	53.4(107)	137	11.7 (133)	123	9.8 (122)	0	* *	0	* *	0	* *	0	* *
0404	176	54.0(104)	158	20.3 (36)	149	20.1 (20)	0	* *	0	* *	0	* *	0	* *
0501	480	61.5 (44)	528	21.6 (31)	471	17.8 (30)	0	* *	0	* *	0	* *	0	* *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in

(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2004 - 2006 Pediatric Nutrition Surveillance (1)
Michigan

Page: 2
Run Date: 5 / 15 / 07

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (4,5)								
	Breastfeeding (5)			Exclusive Breastfeeding (6)			≤2 Hours TV Viewing/Day (7)	Smoking in Household
Clinic	Number (Rank)	% Ever Breastfed (Rank)	Number % Breastfed At Least 6 Months (Rank)	Number % Breastfed At Least 12 Months (Rank)	Number % Breastfed At Least 3 Months (Rank)	Number % Breastfed At Least 6 Months (Rank)	Number (Rank)	% (Rank)
0502	3	* *	3	* *	4	* *	0	* *
0503	182	74.2 (8)	156	28.8 (9)	137	24.8 (10)	0	* *
0504	721	71.8 (11)	637	23.5 (21)	539	15.8 (47)	0	* *
0505	498	64.5 (26)	380	21.6 (31)	319	17.9 (29)	0	* *
0506	222	60.8 (49)	197	15.7 (78)	191	13.1 (70)	0	* *
0650	1,209	70.1 (14)	1,048	18.1 (52)	803	15.4 (52)	0	* *
0651	717	48.5(124)	588	12.4 (124)	443	11.7 (94)	0	* *
0652	464	55.6 (90)	357	17.6 (57)	286	18.5 (25)	0	* *
0653	511	58.9 (64)	447	14.1 (97)	374	12.0 (88)	0	* *
0654	417	62.1 (39)	359	14.2 (95)	294	12.6 (80)	0	* *
0655	513	54.2(101)	439	13.7 (105)	337	12.8 (75)	0	* *
0657	133	63.9 (29)	118	11.0 (142)	115	5.2 (169)	0	* *
0701	498	72.5 (10)	412	32.8 (6)	312	27.2 (7)	0	* *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in

(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2004 - 2006 Pediatric Nutrition Surveillance (1)
Michigan

Page: 3
Run Date: 5 / 15 / 07

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
(4,5)

	Breastfeeding (5)						Exclusive Breastfeeding (6)		≤2 Hours TV Viewing/Day (7)	Smoking in Household		
Clinic	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months	Number	% Breastfed At Least 3 Months	Number	% Breastfed At Least 6 Months	Number	%
	(Rank)		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)	
0702	124	48.4(126)	97	* *	85	* *	0	* *	0	* *	0	* *
0703	274	61.7 (42)	214	13.6 (108)	182	11.0 (103)	0	* *	0	* *	0	* *
0704	98	* *	89	* *	77	* *	0	* *	0	* *	0	* *
0705	202	60.9 (48)	195	23.1 (24)	156	21.2 (14)	0	* *	0	* *	0	* *
0850	1,239	60.0 (56)	1,019	13.2 (116)	788	10.3 (114)	0	* *	0	* *	0	* *
0851	956	51.8(111)	832	13.6 (108)	611	11.3 (98)	0	* *	0	* *	0	* *
0901	2,137	50.5(118)	1,639	13.7 (105)	1,218	9.6 (126)	0	* *	0	* *	0	* *
0902	90	* *	81	* *	63	* *	0	* *	0	* *	0	* *
1051	1,633	77.3 (7)	1,441	31.2 (7)	1,171	26.1 (8)	0	* *	0	* *	0	* *
1059	1	* *	2	* *	3	* *	0	* *	0	* *	0	* *
1060	0	* *	1	* *	2	* *	0	* *	0	* *	0	* *
1101	2,007	44.2(148)	2,056	9.8 (158)	1,511	8.3 (143)	0	* *	0	* *	0	* *
1102	1,115	47.9(133)	913	16.8 (65)	669	16.3 (39)	0	* *	0	* *	0	* *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in

(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2004 - 2006 Pediatric Nutrition Surveillance (1)
Michigan

Page: 4
Run Date: 5 / 15 / 07

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (4,5)														
	Breastfeeding (5)						Exclusive Breastfeeding (6)				≤2 Hours TV Viewing/Day (7)		Smoking in Household	
Clinic	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months	Number	% Breastfed At Least 3 Months	Number	% Breastfed At Least 6 Months	Number	%	Number	%
		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)
1106	137	55.5 (91)	124	12.9 (118)	88	* *	0	* *	0	* *	0	* *	0	* *
1250	1,163	55.5 (91)	912	15.4 (83)	783	13.8 (62)	0	* *	0	* *	0	* *	0	* *
1251	797	56.0 (85)	812	11.0 (142)	621	8.1 (146)	0	* *	0	* *	0	* *	0	* *
1252	991	52.6(109)	761	16.6 (67)	679	13.8 (62)	0	* *	0	* *	0	* *	0	* *
1253	919	42.3(155)	740	7.8 (171)	535	7.1 (159)	0	* *	0	* *	0	* *	0	* *
1350	2,907	43.9(149)	2,258	9.6 (161)	1,776	8.0 (148)	0	* *	0	* *	0	* *	0	* *
1351	219	59.4 (61)	203	19.2 (40)	164	14.0 (60)	0	* *	0	* *	0	* *	0	* *
1352	551	37.2(165)	423	9.7 (159)	353	8.2 (145)	0	* *	0	* *	0	* *	0	* *
1401	314	73.2 (9)	272	26.5 (12)	228	18.9 (24)	0	* *	0	* *	0	* *	0	* *
1402	209	82.3 (4)	184	27.2 (11)	146	15.8 (47)	0	* *	0	* *	0	* *	0	* *
1403	34	* *	31	* *	31	* *	0	* *	0	* *	0	* *	0	* *
1404	25	* *	29	* *	26	* *	0	* *	0	* *	0	* *	0	* *
1501	1,334	57.7 (72)	1,081	19.0 (42)	891	16.7 (36)	0	* *	0	* *	0	* *	0	* *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in

(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2004 - 2006 Pediatric Nutrition Surveillance (1)
Michigan

Page: 5
Run Date: 5 / 15 / 07

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
(4,5)

	Breastfeeding (5)						Exclusive Breastfeeding (6)				≤2 Hours TV Viewing/Day (7)	Smoking in Household		
Clinic	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months	Number	% Breastfed At Least 3 Months	Number	% Breastfed At Least 6 Months	Number	%	Number	%
	(Rank)		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)	
1502	156	57.7 (72)	130	12.3 (125)	103	7.8 (149)	0	* *	0	* *	0	* *	0	* *
1503	1,335	58.6 (67)	1,147	18.7 (46)	961	14.2 (58)	0	* *	0	* *	0	* *	0	* *
1504	2,904	63.5 (32)	2,407	19.9 (37)	2,011	16.3 (39)	0	* *	0	* *	0	* *	0	* *
1505	819	64.8 (24)	741	24.0 (20)	614	16.1 (43)	0	* *	0	* *	0	* *	0	* *
1506	621	54.1(103)	516	14.3 (94)	418	11.0 (103)	0	* *	0	* *	0	* *	0	* *
1507	723	59.8 (57)	626	19.0 (42)	502	15.5 (50)	0	* *	0	* *	0	* *	0	* *
1508	272	79.0 (6)	268	24.3 (19)	249	17.7 (32)	0	* *	0	* *	0	* *	0	* *
1509	753	61.0 (47)	646	19.8 (38)	559	18.1 (28)	0	* *	0	* *	0	* *	0	* *
1510	263	54.4 (99)	224	15.6 (81)	191	11.5 (96)	0	* *	0	* *	0	* *	0	* *
1511	654	58.0 (69)	539	17.4 (59)	406	14.3 (57)	0	* *	0	* *	0	* *	0	* *
1512	359	57.9 (70)	298	15.1 (86)	224	11.2 (100)	0	* *	0	* *	0	* *	0	* *
1701	749	64.0 (28)	560	21.8 (30)	453	15.9 (46)	0	* *	0	* *	0	* *	0	* *
1712	136	57.4 (77)	113	20.4 (35)	114	12.3 (84)	0	* *	0	* *	0	* *	0	* *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in

(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2004 - 2006 Pediatric Nutrition Surveillance (1)
Michigan

Page: 6
Run Date: 5 / 15 / 07

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (4,5)														
	Breastfeeding (5)						Exclusive Breastfeeding (6)				≤2 Hours TV Viewing/Day (7)		Smoking in Household	
Clinic	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months	Number	% Breastfed At Least 3 Months	Number	% Breastfed At Least 6 Months	Number	%	Number	%
		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)
1950	813	55.8 (87)	660	15.6 (81)	528	12.1 (87)	0	* *	0	* *	0	* *	0	* *
1952	200	54.0(104)	141	16.3 (73)	124	18.5 (25)	0	* *	0	* *	0	* *	0	* *
1953	442	54.3(100)	341	16.7 (66)	272	14.7 (55)	0	* *	0	* *	0	* *	0	* *
1954	462	58.7 (65)	373	18.0 (53)	292	12.7 (77)	0	* *	0	* *	0	* *	0	* *
1955	232	56.5 (80)	177	11.9 (130)	137	10.2 (115)	0	* *	0	* *	0	* *	0	* *
1956	8	* *	10	* *	8	* *	0	* *	0	* *	0	* *	0	* *
1957	470	61.9 (41)	407	14.7 (89)	303	12.9 (73)	0	* *	0	* *	0	* *	0	* *
1959	250	60.8 (49)	224	16.1 (75)	169	10.7 (107)	0	* *	0	* *	0	* *	0	* *
1963	243	64.2 (27)	202	14.4 (92)	160	10.0 (121)	0	* *	0	* *	0	* *	0	* *
2150	810	59.5 (60)	680	15.7 (78)	592	12.2 (86)	0	* *	0	* *	0	* *	0	* *
2151	361	48.2(131)	275	10.9 (144)	234	7.7 (152)	0	* *	0	* *	0	* *	0	* *
2251	622	57.7 (72)	538	17.8 (54)	421	13.5 (66)	0	* *	0	* *	0	* *	0	* *
2252	209	56.5 (80)	185	13.5 (111)	147	12.9 (73)	0	* *	0	* *	0	* *	0	* *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in

(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2004 - 2006 Pediatric Nutrition Surveillance (1)
Michigan

Page: 7
Run Date: 5 / 15 / 07

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
(4,5)

	Breastfeeding (5)						Exclusive Breastfeeding (6)				≤2 Hours TV Viewing/Day (7)	Smoking in Household		
Clinic	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months	Number	% Breastfed At Least 3 Months	Number	% Breastfed At Least 6 Months	Number	%	Number	%
		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)
2551	4,380	35.0(169)	3,202	6.8 (175)	2,310	5.4 (168)	0	* *	0	* *	0	* *	0	* *
2552	1,084	41.8(156)	823	9.7 (159)	583	9.8 (122)	0	* *	0	* *	0	* *	0	* *
2554	133	41.4(158)	102	6.9 (174)	87	* *	0	* *	0	* *	0	* *	0	* *
2555	4,706	48.1(132)	3,453	10.6 (149)	2,494	8.4 (141)	0	* *	0	* *	0	* *	0	* *
3001	1,197	36.4(167)	888	8.6 (165)	700	7.4 (157)	0	* *	0	* *	0	* *	0	* *
3002	251	35.1(168)	195	8.2 (168)	163	5.5 (167)	0	* *	0	* *	0	* *	0	* *
3003	26	* *	24	* *	17	* *	0	* *	0	* *	0	* *	0	* *
3004	259	80.7 (5)	217	42.4 (4)	189	30.7 (5)	0	* *	0	* *	0	* *	0	* *
3007	44	* *	38	* *	34	* *	0	* *	0	* *	0	* *	0	* *
3009	10	* *	11	* *	7	* *	0	* *	0	* *	0	* *	0	* *
3010	1	* *	2	* *	1	* *	0	* *	0	* *	0	* *	0	* *
3201	720	45.6(141)	569	11.1 (140)	426	10.1 (117)	0	* *	0	* *	0	* *	0	* *
3202	10	* *	12	* *	16	* *	0	* *	0	* *	0	* *	0	* *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in

(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2004 - 2006 Pediatric Nutrition Surveillance (1)
Michigan

Page: 8
Run Date: 5 / 15 / 07

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
(4,5)

	Breastfeeding (5)						Exclusive Breastfeeding (6)				≤2 Hours TV Viewing/Day (7)	Smoking in Household		
Clinic	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months	Number	% Breastfed At Least 3 Months	Number	% Breastfed At Least 6 Months	Number	%	Number	%
		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)
3301	3,825	59.1 (63)	3,857	17.3 (61)	2,972	13.8 (62)	0	* *	0	* *	0	* *	0	* *
3302	439	56.0 (85)	465	14.4 (92)	378	13.8 (62)	0	* *	0	* *	0	* *	0	* *
3303	87	* *	82	* *	58	* *	0	* *	0	* *	0	* *	0	* *
3304	170	94.1 (1)	200	52.0 (2)	162	43.2 (1)	0	* *	0	* *	0	* *	0	* *
3305	116	64.7 (25)	127	22.8 (25)	109	22.0 (12)	0	* *	0	* *	0	* *	0	* *
3306	109	61.5 (44)	134	9.0 (163)	113	6.2 (165)	0	* *	0	* *	0	* *	0	* *
3308	60	* *	61	* *	52	* *	0	* *	0	* *	0	* *	0	* *
3309	42	* *	45	* *	32	* *	0	* *	0	* *	0	* *	0	* *
3311	14	* *	12	* *	8	* *	0	* *	0	* *	0	* *	0	* *
3550	108	45.4(142)	75	* *	73	* *	0	* *	0	* *	0	* *	0	* *
3551	41	* *	26	* *	25	* *	0	* *	0	* *	0	* *	0	* *
3552	46	* *	46	* *	39	* *	0	* *	0	* *	0	* *	0	* *
3650	27	* *	26	* *	24	* *	0	* *	0	* *	0	* *	0	* *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in

(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2004 - 2006 Pediatric Nutrition Surveillance (1)
Michigan

Page: 9
Run Date: 5 / 15 / 07

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
(4,5)

	Breastfeeding (5)			Exclusive Breastfeeding (6)			≤2 Hours TV Viewing/Day (7)	Smoking in Household
Clinic	Number (Rank)	% Ever Breastfed (Rank)	Number At Least 6 Months (Rank)	% Breastfed At Least 12 Months (Rank)	Number At Least 3 Months (Rank)	% Breastfed At Least 6 Months (Rank)	Number (Rank)	% (Rank)
3701	836	52.8(108)	685	13.7 (105)	526	12.7 (77)	0	* *
3702	281	51.2(116)	219	8.2 (168)	142	4.9 (170)	0	* *
3703	136	47.8(135)	104	12.5 (123)	66	* *	0	* *
3704	120	50.8(117)	106	15.1 (86)	72	* *	0	* *
3805	3,671	50.4(120)	3,116	11.6 (135)	2,258	8.4 (141)	0	* *
3950	2,383	55.8 (87)	1,865	17.2 (62)	1,391	15.5 (50)	0	* *
3952	80	* *	77	* *	57	* *	0	* *
3953	105	67.6 (17)	72	* *	55	* *	0	* *
3954	227	54.2(101)	183	21.9 (29)	136	14.7 (55)	0	* *
4001	1,147	55.1 (94)	908	13.0 (117)	747	10.6 (109)	0	* *
4002	941	57.8 (71)	760	13.9 (100)	591	12.0 (88)	0	* *
4103	51	* *	47	* *	44	* *	0	* *
4106	1,826	51.6(113)	1,601	13.3 (115)	1,270	10.6 (109)	0	* *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in

(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2004 - 2006 Pediatric Nutrition Surveillance (1)
Michigan

Page: 10
Run Date: 5 / 15 / 07

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
(4,5)

	Breastfeeding (5)						Exclusive Breastfeeding (6)				≤2 Hours TV Viewing/Day (7)	Smoking in Household		
Clinic	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months	Number	% Breastfed At Least 3 Months	Number	% Breastfed At Least 6 Months	Number	%	Number	%
	(Rank)		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)	
4107	3,323	56.5 (80)	3,044	16.4 (71)	2,365	12.8 (75)	0	* *	0	* *	0	* *	0	* *
4109	1,260	58.7 (65)	1,024	15.2 (85)	831	9.7 (124)	0	* *	0	* *	0	* *	0	* *
4110	1,841	59.8 (57)	1,623	18.9 (44)	1,311	15.4 (52)	0	* *	0	* *	0	* *	0	* *
4111	65	* *	50	* *	35	* *	0	* *	0	* *	0	* *	0	* *
4112	833	90.6 (2)	716	19.1 (41)	638	16.8 (35)	0	* *	0	* *	0	* *	0	* *
4113	663	71.8 (11)	677	20.8 (34)	600	16.5 (37)	0	* *	0	* *	0	* *	0	* *
4120	2,328	60.2 (53)	2,003	16.4 (71)	1,567	13.3 (68)	0	* *	0	* *	0	* *	0	* *
4160	2,201	55.3 (93)	1,854	15.8 (77)	1,389	12.7 (77)	0	* *	0	* *	0	* *	0	* *
4351	830	56.6 (79)	676	15.7 (78)	547	10.1 (117)	0	* *	0	* *	0	* *	0	* *
4352	256	48.4(126)	205	11.7 (133)	167	7.8 (149)	0	* *	0	* *	0	* *	0	* *
4353	607	62.9 (35)	492	16.5 (69)	429	11.7 (94)	0	* *	0	* *	0	* *	0	* *
4354	925	63.1 (33)	772	17.0 (64)	587	12.3 (84)	0	* *	0	* *	0	* *	0	* *
4355	418	62.7 (37)	325	16.0 (76)	298	16.1 (43)	0	* *	0	* *	0	* *	0	* *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in

(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2004 - 2006 Pediatric Nutrition Surveillance (1)
Michigan

Page: 11
Run Date: 5 / 15 / 07

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
(4,5)

	Breastfeeding (5)			Exclusive Breastfeeding (6)			≤2 Hours TV Viewing/Day (7)	Smoking in Household
Clinic	Number (Rank)	% Ever Breastfed (Rank)	Number (Rank)	% Breastfed At Least 6 Months (Rank)	Number (Rank)	% Breastfed At Least 12 Months (Rank)	Number (Rank)	% (Rank)
4356	654	59.2 (62)	526	17.7 (55)	461	9.3 (132)	0	* *
4357	19	* *	18	* *	16	* *	0	* *
4358	300	62.0 (40)	287	28.9 (8)	266	21.1 (15)	0	* *
4359	423	62.2 (38)	305	18.7 (46)	289	15.6 (49)	0	* *
4360	220	65.5 (22)	166	18.7 (46)	142	11.3 (98)	0	* *
4361	287	60.3 (52)	216	11.1 (140)	188	9.0 (133)	0	* *
4362	434	71.2 (13)	350	17.7 (55)	285	18.2 (27)	0	* *
4363	1,016	63.0 (34)	808	14.2 (95)	639	8.6 (138)	0	* *
4401	1,376	51.4(115)	1,082	13.9 (100)	888	9.7 (124)	0	* *
4501	1,218	58.3 (68)	1,073	10.2 (154)	869	8.5 (140)	0	* *
4601	1,735	52.2(110)	1,461	12.3 (125)	1,043	10.6 (109)	0	* *
4701	1,296	66.4 (20)	1,022	17.4 (59)	745	13.2 (69)	0	* *
5050	1,810	44.3(145)	1,352	11.4 (136)	915	9.4 (131)	0	* *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in

(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2004 - 2006 Pediatric Nutrition Surveillance (1)
Michigan

Page: 12
Run Date: 5 / 15 / 07

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
(4,5)

	Breastfeeding (5)						Exclusive Breastfeeding (6)				≤2 Hours TV Viewing/Day (7)	Smoking in Household		
Clinic	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months	Number	% Breastfed At Least 3 Months	Number	% Breastfed At Least 6 Months	Number	%	Number	%
		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)
5051	4,608	48.3(129)	3,557	12.9 (118)	2,563	9.6 (126)	0	* * *	0	* * *	0	* * *	0	* * *
5052	2,728	40.9(161)	2,069	10.5 (151)	1,439	8.1 (146)	0	* * *	0	* * *	0	* * *	0	* * *
5054	312	62.8 (36)	240	24.6 (17)	185	16.2 (41)	0	* * *	0	* * *	0	* * *	0	* * *
5202	380	65.3 (23)	311	23.5 (21)	246	19.5 (21)	0	* * *	0	* * *	0	* * *	0	* * *
5203	230	56.5 (80)	192	17.2 (62)	164	13.4 (67)	0	* * *	0	* * *	0	* * *	0	* * *
5204	168	68.5 (16)	133	28.6 (10)	115	19.1 (23)	0	* * *	0	* * *	0	* * *	0	* * *
5205	281	60.1 (55)	220	25.0 (16)	180	20.6 (18)	0	* * *	0	* * *	0	* * *	0	* * *
5601	1,257	61.7 (42)	1,095	15.4 (83)	824	11.5 (96)	0	* * *	0	* * *	0	* * *	0	* * *
5801	2,331	50.5(118)	1,886	13.4 (112)	1,340	10.1 (117)	0	* * *	0	* * *	0	* * *	0	* * *
5803	17	* * *	11	* * *	10	* * *	0	* * *	0	* * *	0	* * *	0	* * *
6103	4,914	43.5(150)	3,217	12.0 (129)	3,078	7.7 (152)	0	* * *	0	* * *	0	* * *	0	* * *
6301	5,329	47.7(137)	4,092	14.6 (90)	2,915	12.6 (80)	0	* * *	0	* * *	0	* * *	0	* * *
6312	3,133	54.7 (98)	2,522	16.5 (69)	1,794	13.0 (71)	0	* * *	0	* * *	0	* * *	0	* * *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in

(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2004 - 2006 Pediatric Nutrition Surveillance (1)
Michigan

Page: 13
Run Date: 5 / 15 / 07

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
(4,5)

	Breastfeeding (5)						Exclusive Breastfeeding (6)				≤2 Hours TV Viewing/Day (7)	Smoking in Household		
Clinic	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months	Number	% Breastfed At Least 3 Months	Number	% Breastfed At Least 6 Months	Number	%	Number	%
	(Rank)		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)	
6316	1,333	57.5 (76)	1,110	14.0 (98)	798	11.2 (100)	0	* *	0	* *	0	* *	0	* *
6320	329	60.2 (53)	289	26.3 (14)	211	28.0 (6)	0	* *	0	* *	0	* *	0	* *
6325	189	50.3(121)	166	13.9 (100)	120	14.2 (58)	0	* *	0	* *	0	* *	0	* *
6330	319	40.4(162)	254	10.6 (149)	166	12.0 (88)	0	* *	0	* *	0	* *	0	* *
6335	724	50.0(122)	613	21.0 (33)	465	21.5 (13)	0	* *	0	* *	0	* *	0	* *
6340	639	54.0(104)	496	18.8 (45)	345	14.8 (54)	0	* *	0	* *	0	* *	0	* *
7304	339	47.8(135)	248	10.9 (144)	227	6.6 (162)	0	* *	0	* *	0	* *	0	* *
7306	3,147	44.6(143)	2,046	6.8 (175)	1,889	4.5 (172)	0	* *	0	* *	0	* *	0	* *
7308	251	63.7 (30)	194	13.4 (112)	166	9.6 (126)	0	* *	0	* *	0	* *	0	* *
7401	2,666	44.3(145)	2,036	10.0 (157)	1,525	8.3 (143)	0	* *	0	* *	0	* *	0	* *
7402	384	48.4(126)	315	10.5 (151)	229	6.6 (162)	0	* *	0	* *	0	* *	0	* *
7403	118	45.8(140)	94	* *	72	* *	0	* *	0	* *	0	* *	0	* *
7404	157	51.6(113)	134	12.7 (121)	101	11.9 (92)	0	* *	0	* *	0	* *	0	* *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in

(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2004 - 2006 Pediatric Nutrition Surveillance (1)
Michigan

Page: 14
Run Date: 5 / 15 / 07

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic														
(4,5)														
	Breastfeeding (5)						Exclusive Breastfeeding (6)				≤2 Hours TV Viewing/Day (7)		Smoking in Household	
Clinic	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months	Number	% Breastfed At Least 3 Months	Number	% Breastfed At Least 6 Months	Number	%	Number	%
		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)
7601	857	56.5 (80)	729	12.2 (127)	578	10.2 (115)	0	* *	0	* *	0	* *	0	* *
7850	1,251	55.7 (89)	948	14.9 (88)	760	10.5 (112)	0	* *	0	* *	0	* *	0	* *
7851	262	67.6 (17)	249	12.9 (118)	218	7.8 (149)	0	* *	0	* *	0	* *	0	* *
7852	143	63.6 (31)	135	16.3 (73)	102	8.8 (135)	0	* *	0	* *	0	* *	0	* *
8150	127	66.9 (19)	90	* *	51	* *	0	* *	0	* *	0	* *	0	* *
8151	3,051	61.3 (46)	2,264	22.8 (25)	1,694	17.4 (33)	0	* *	0	* *	0	* *	0	* *
8152	531	54.8 (97)	379	7.9 (170)	261	6.9 (160)	0	* *	0	* *	0	* *	0	* *
8153	44	* *	37	* *	28	* *	0	* *	0	* *	0	* *	0	* *
8154	462	88.7 (3)	403	61.5 (1)	343	43.1 (2)	0	* *	0	* *	0	* *	0	* *
8155	90	* *	68	* *	56	* *	0	* *	0	* *	0	* *	0	* *
8156	54	* *	44	* *	40	* *	0	* *	0	* *	0	* *	0	* *
8157	13	* *	10	* *	11	* *	0	* *	0	* *	0	* *	0	* *
8201	3,030	47.2(139)	2,152	12.6 (122)	1,480	10.1 (117)	0	* *	0	* *	0	* *	0	* *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in

(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2004 - 2006 Pediatric Nutrition Surveillance (1)
Michigan

Page: 15
Run Date: 5 / 15 / 07

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (4,5)														
	Breastfeeding (5)						Exclusive Breastfeeding (6)				≤2 Hours TV Viewing/Day (7)		Smoking in Household	
Clinic	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months	Number	% Breastfed At Least 3 Months	Number	% Breastfed At Least 6 Months	Number	%	Number	%
		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)
8202	3,507	68.7 (15)	2,507	43.1 (3)	2,444	40.3 (3)	0	* *	0	* *	0	* *	0	* *
8253	374	47.9(133)	330	14.5 (91)	254	16.1 (43)	0	* *	0	* *	0	* *	0	* *
8254	432	34.5(170)	312	12.2 (127)	291	11.0 (103)	0	* *	0	* *	0	* *	0	* *
8255	3,768	43.1(152)	2,300	13.8 (104)	1,812	11.8 (93)	0	* *	0	* *	0	* *	0	* *
8257	709	48.7(123)	418	19.4 (39)	393	17.0 (34)	0	* *	0	* *	0	* *	0	* *
8258	2,058	41.5(157)	1,195	10.8 (146)	1,153	7.6 (154)	0	* *	0	* *	0	* *	0	* *
8901	1,228	30.0(175)	874	13.4 (112)	618	10.4 (113)	0	* *	0	* *	0	* *	0	* *
8902	642	37.1(166)	478	18.2 (51)	358	16.5 (37)	0	* *	0	* *	0	* *	0	* *
8903	1,356	29.0(176)	1,021	13.9 (100)	712	11.0 (103)	0	* *	0	* *	0	* *	0	* *
8904	885	47.6(138)	719	26.4 (13)	574	22.1 (11)	0	* *	0	* *	0	* *	0	* *
8905	2,301	37.4(164)	1,567	22.3 (28)	1,132	20.2 (19)	0	* *	0	* *	0	* *	0	* *
9001	2	* *	3	* *	11	* *	0	* *	0	* *	0	* *	0	* *
9002	7	* *	12	* *	15	* *	0	* *	0	* *	0	* *	0	* *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in

(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2004 - 2006 Pediatric Nutrition Surveillance (1)
Michigan

Page: 16
Run Date: 5 / 15 / 07

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
(4,5)

	Breastfeeding (5)						Exclusive Breastfeeding (6)				≤2 Hours TV Viewing/Day (7)		Smoking in Household	
Clinic	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months	Number	% Breastfed At Least 3 Months	Number	% Breastfed At Least 6 Months	Number	%	Number	%
	(Rank)		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)	
9004	393	25.7(180)	422	10.7 (148)	334	12.0 (88)	0	* * *	0	* * *	0	* * *	0	* * *
9005	4	* * *	5	* * *	6	* * *	0	* * *	0	* * *	0	* * *	0	* * *
9006	8	* * *	10	* * *	14	* * *	0	* * *	0	* * *	0	* * *	0	* * *
9008	1	* * *	1	* * *	4	* * *	0	* * *	0	* * *	0	* * *	0	* * *
9009	13	* * *	22	* * *	16	* * *	0	* * *	0	* * *	0	* * *	0	* * *
9010	254	21.3(182)	258	10.5 (151)	206	10.7 (107)	0	* * *	0	* * *	0	* * *	0	* * *
9011	5	* * *	3	* * *	8	* * *	0	* * *	0	* * *	0	* * *	0	* * *
9012	7	* * *	12	* * *	6	* * *	0	* * *	0	* * *	0	* * *	0	* * *
9014	7	* * *	13	* * *	8	* * *	0	* * *	0	* * *	0	* * *	0	* * *
9018	1	* * *	5	* * *	7	* * *	0	* * *	0	* * *	0	* * *	0	* * *
9020	4	* * *	10	* * *	10	* * *	0	* * *	0	* * *	0	* * *	0	* * *
9022	3	* * *	5	* * *	10	* * *	0	* * *	0	* * *	0	* * *	0	* * *
9023	19	* * *	21	* * *	25	* * *	0	* * *	0	* * *	0	* * *	0	* * *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in

(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2004 - 2006 Pediatric Nutrition Surveillance (1)
Michigan

Page: 17
Run Date: 5 / 15 / 07

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (4,5)								
	Breastfeeding (5)			Exclusive Breastfeeding (6)			≤2 Hours TV Viewing/Day (7)	Smoking in Household
Clinic	Number (Rank)	% Ever Breastfed (Rank)	Number % Breastfed At Least 6 Months (Rank)	Number % Breastfed At Least 12 Months (Rank)	Number % Breastfed At Least 3 Months (Rank)	Number % Breastfed At Least 6 Months (Rank)	Number (Rank)	% (Rank)
9024	7	* *	13	* *	12	* *	0	* *
9035	8	* *	18	* *	14	* *	0	* *
9039	19	* *	27	* *	22	* *	0	* *
9040	310	27.1(179)	352	10.2 (154)	268	9.0 (133)	0	* *
9045	1,470	40.4(162)	1,077	18.3 (49)	946	17.8 (30)	0	* *
9050	2,207	48.3(129)	1,791	24.5 (18)	1,337	21.1 (15)	0	* *
9051	2,678	54.9 (96)	1,560	23.3 (23)	1,731	19.5 (21)	0	* *
9053	3,089	32.2(172)	1,909	13.6 (108)	1,692	12.5 (82)	0	* *
9054	788	34.1(171)	561	7.0 (173)	372	7.5 (156)	0	* *
9058	573	55.1 (94)	481	22.5 (27)	399	20.8 (17)	0	* *
9066	2,922	28.0(178)	1,513	11.4 (136)	1,492	9.5 (130)	0	* *
9068	1,671	31.1(173)	1,174	8.6 (165)	930	7.2 (158)	0	* *
9069	2,461	41.3(160)	1,911	11.3 (139)	1,281	9.6 (126)	0	* *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in

(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2004 - 2006 Pediatric Nutrition Surveillance (1)
Michigan

Page: 18
Run Date: 5 / 15 / 07

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic														
(4,5)														
	Breastfeeding (5)						Exclusive Breastfeeding (6)				≤2 Hours TV Viewing/Day (7)		Smoking in Household	
Clinic	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months	Number	% Breastfed At Least 3 Months	Number	% Breastfed At Least 6 Months	Number	%	Number	%
		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)
9070	2,336	30.4(174)	1,940	9.3 (162)	1,412	8.7 (136)	0	* *	0	* *	0	* *	0	* *
9072	1,876	28.1(177)	956	9.0 (163)	996	6.8 (161)	0	* *	0	* *	0	* *	0	* *
9074	1,565	43.1(152)	1,247	11.9 (130)	848	8.6 (138)	0	* *	0	* *	0	* *	0	* *
9079	1,516	43.5(150)	1,178	11.9 (130)	771	11.2 (100)	0	* *	0	* *	0	* *	0	* *
9080	2,289	56.9 (78)	1,965	33.0 (5)	1,723	33.0 (4)	0	* *	0	* *	0	* *	0	* *
9151	15	* *	8	* *	6	* *	0	* *	0	* *	0	* *	0	* *
Unknown Clinics	614	24.9(181)	435	8.3 (167)	286	4.9 (170)	0	* *	0	* *	0	* *	0	* *
Michigan	193,062	50.8	151,645	15.7	119,710	13.1	0	*	0	*	0	*	0	*

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in

(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.